



Guiding Principles and Elements of Recovery-Oriented Systems of Care:

What do we know from the research?

August 2009



U.S. DEPARTMENT OF HEALTH AND HUMAN SERVICES
Substance Abuse and Mental Health Services Administration
Center for Substance Abuse Treatment
www.samhsa.gov

Guiding Principles and Elements of Recovery-Oriented Systems of Care:

What do we know from the research?

U.S. Department of Health and Human Services
Substance Abuse and Mental Health Services Administration
Center for Substance Abuse Treatment

Acknowledgments

This publication was prepared for the Substance Abuse and Mental Health Services Administration (SAMHSA) by Cori Kautz Sheedy, M.A., of Abt Associates Inc., under the direction of Melanie Whitter, Abt Associates Inc., under contract number 270-03-9000, with SAMHSA, U.S. Department of Health and Human Services (HHS). Shannon B. Taitt, M.P.A., served as the Government Project Officer.

Disclaimer

The views, opinions, and content of this publication are those of the authors and do not necessarily reflect the views, opinions, or policies of SAMHSA or HHS.

Public Domain Notice

All material appearing in this report is in the public domain and may be reproduced or copied without permission from SAMHSA. Citation of the source is appreciated. However, this publication may *not* be reproduced or distributed for a fee without the specific, written authorization of the Office of Communications, SAMHSA, HHS.

Electronic Access and Copies of Publication

This publication may be downloaded at <http://www.samhsa.gov/shin> or <http://pfr.samhsa.gov/ros.html>. Or, please call SAMHSA's Health Information Network at 1-877-SAMHSA-7 (1-877-726-4727) (English and Español).

Recommended Citation

Sheedy C. K., and Whitter M., *Guiding Principles and Elements of Recovery-Oriented Systems of Care: What Do We Know From the Research?* HHS Publication No. (SMA) 09-4439. Rockville, MD: Center for Substance Abuse Treatment, Substance Abuse and Mental Health Services Administration, 2009.

Originating Office

Office of Program Analysis and Coordination, Center for Substance Abuse Treatment, Substance Abuse and Mental Health Services Administration, 1 Choke Cherry Road, Rockville, MD 20857.

HHS Publication No. (SMA) 09-4439

Table of Contents

1. Background	1
2. Purpose Statement	4
3. Methodology	5
4. Research Supporting the Conceptual Framework of Recovery-Oriented Systems of Care.....	6
5. Research Supporting the Principles and Systems of Care Elements	13
6. Research Supporting the Principles of Recovery.....	15
7. Research Supporting the Systems of Care Elements	22
8. Research Supporting the Implementation of Recovery-Oriented Services and Systems of Care.....	34
9. Conclusion	39
References	41

1. Background

The concept of recovery lies at the core of the Substance Abuse and Mental Health Services Administration's (SAMHSA's) mission, and fostering the development of recovery-oriented systems of care and services is a Center for Substance Abuse Treatment (CSAT) priority. In support of that commitment, in 2005, SAMHSA's CSAT convened a National Summit on Recovery. Participants at the Summit represented a broad group of stakeholders, policymakers, advocates, consumers, clinicians, and administrators from diverse ethnic and professional backgrounds. Although the substance use problems and disorders treatment and recovery field has discussed and lived recovery for decades, the Summit represented the first broad-based national effort to reach a definition of recovery and a common understanding of the guiding principles of recovery and the elements of recovery-oriented systems of care.

Through a multistage process, key stakeholders formulated guiding principles of recovery and key elements of recovery-oriented systems of care. Summit participants then further refined the guiding principles and key elements in response to two questions: 1) What principles of recovery should guide the field in the future? and 2) What ideas could help make the field more recovery oriented?

A working definition of recovery, 12 guiding principles of recovery, and 17 elements of recovery-oriented systems of care emerged from the Summit process; these are subsequently defined in this paper and in the

*National Summit on Recovery: Conference Report.*¹ These principles and elements can now provide a philosophical and conceptual framework to guide SAMHSA/CSAT and other stakeholder groups and offer a shared language for dialog.

Summit participants agreed on the following working definition of recovery:

Recovery from alcohol and drug problems is a process of change through which an individual achieves abstinence and improved health, wellness, and quality of life.

The guiding principles that emerged from the Summit are broad and overarching. They are intended to give general direction to SAMHSA/CSAT and other stakeholder groups as the treatment and recovery field moves toward operationalizing recovery-oriented systems of care and developing core measures, promising approaches, and evidence-based practices. The principles also helped Summit participants define the recovery-oriented elements and guided recommendations for the field.

Following are the 12 guiding principles identified by participants (defined in this paper):

- There are many pathways to recovery.
- Recovery is self-directed and empowering.

- Recovery involves a personal recognition of the need for change and transformation.
- Recovery is holistic.
- Recovery has cultural dimensions.
- Recovery exists on a continuum of improved health and wellness.
- Recovery emerges from hope and gratitude.
- Recovery involves a process of healing and self-redefinition.
- Recovery involves addressing discrimination and transcending shame and stigma.
- Recovery is supported by peers and allies.
- Recovery involves (re)joining and (re)building a life in the community.
- Recovery is a reality.

Participants at the Summit agreed that recovery-oriented systems of care are as complex and dynamic as the process of recovery itself. They are designed to support individuals seeking to overcome substance use problems and disorders across their lifespan. Participants at the Summit declared, “There will be no wrong door to recovery” and also recognized that recovery-oriented systems of care need to provide “genuine, free and independent choice” among an array of treatment and recovery support options. Services should optimally be provided in flexible, unbundled packages that evolve over time to meet the changing needs of recovering individuals. Individuals should also be able to access a comprehensive array of services that are fully coordinated to provide support to individuals

throughout their unique journeys to sustained recovery.


Summit participants identified the following 17 elements of recovery-oriented systems of care and services (defined in this paper):

- Person-centered;
- Inclusive of family and other ally involvement;
- Individualized and comprehensive services across the lifespan;
- Systems anchored in the community;
- Continuity of care;
- Partnership-consultant relationships;
- Strength-based;
- Culturally responsive;
- Responsiveness to personal belief systems;
- Commitment to peer recovery support services;
- Inclusion of the voices and experiences of recovering individuals and their families;
- Integrated services;
- System-wide education and training;
- Ongoing monitoring and outreach;
- Outcomes driven;
- Research based; and
- Adequately and flexibly financed.

Work conducted after the Summit defined recovery-oriented systems of care as networks of organizations, agencies, and community members that coordinate a wide spectrum of services to prevent, intervene in, and treat

substance use problems and disorders. Figure 1 illustrates the components of the recovery-oriented systems of care framework.

Figure 1: Conceptual Framework of a Recovery-Oriented System of Care



2. Purpose Statement

The purpose of this white paper is to review the research related to the 12 guiding principles of recovery and the 17 elements of recovery-oriented systems of care developed through the National Summit on Recovery. It also offers an appraisal of scientific literature discussing the recovery-oriented systems of care conceptual framework and literature on recovery-oriented service and systems implementation.

Policymakers, providers, practitioners, researchers, recovery support staff, and others interested in the concepts of recovery and recovery-oriented systems of care and services frequently seek data to inform policy development, planning, and program and systems development. This white paper has been prepared as a resource to those seeking information on the research related to recovery.

States, communities, and organizations across the nation are developing and implementing recovery-oriented services and systems. In this paper, they will find evidence that supports and validates services and systems improvements based on recovery-oriented approaches.

Finally, this paper establishes a baseline of existing research upon which the treatment and recovery field can continue to build an understanding of recovery and recovery-oriented systems of care. This paper also identifies areas in the recovery research where additional data are needed.

3. Methodology

Studies included in this paper were identified by searches of electronic bibliographic databases (PsychINFO, PubMed) as well as citations in published studies. Using keywords and established selection criteria related to the principles and systems elements, we conducted a computerized search of health, addictions, financial, and trade journals and newsletters. Many

relevant studies were identified. This paper includes a review of research related to recovery-oriented systems of care and services for addiction published in peer-reviewed journals, books, and government publications within the past 20 years, with a focus on the last 10 years.

4. Research Supporting the Conceptual Framework of Recovery-Oriented Systems of Care

Recovery has been called the “organizing construct” for the addictions field.² Recently, a conceptual framework that describes and coordinates the delivery of care for individuals with substance use problems and disorders has begun to emerge. Recovery-oriented systems of care are networks of organizations, agencies, and community members that coordinate a wide spectrum of services to prevent, intervene in, and treat substance use problems and disorders. Although States and communities are implementing a variety of services and activities to create recovery-oriented systems, there is minimal research in peer-reviewed journals that examines the framework and the effectiveness and outcomes of this framework. Research is beginning to emerge within the mental health and addictions field related to the recovery-oriented systems of care framework, but the literature is scant. Systems of care research in the addictions field, although conducted within a treatment construct, provides helpful systems information. This section outlines research from the addictions and mental health fields supporting the recovery-oriented systems of care framework, supplemented by research on a systems of care approach within a treatment construct.

In a 2005 article in the *Psychiatric Rehabilitation Journal*, O’Connell, Tondora, Croog, Evans, and Davidson³ conducted a comprehensive review of the literature on mental illness and addictions recovery that

identified the elements of a recovery-oriented environment. According to the authors, a recovery-oriented environment is one that:

- Encourages individuality;
- Promotes accurate and positive portrayals of psychiatric disability while fighting discrimination;
- Focuses on strengths;
- Uses a language of hope and possibility;
- Offers a variety of options for treatment, rehabilitation, and support;
- Supports risk-taking, even when failure is a possibility;
- Actively involves service users, family members, and other natural supports in the development and implementation of programs and services;
- Encourages user participation in advocacy activities;
- Helps develop connections with communities; and
- Helps people develop valued social roles, interests and hobbies, and other meaningful activities.

Gagne, White, and Anthony in a 2007 article in the *Psychiatric Rehabilitation Journal* describe the recovery vision and the values of recovery-oriented care that intersect the addiction and mental health fields:⁴

- Recovery is a personal and individualized process of

growth that unfolds along a continuum, and there are multiple pathways to recovery.

- People in recovery are active agents of change in their lives and not passive recipients of services.
- People in recovery from mental illness and/or addiction disorders often note the important role of family and peer support in making the difference in their recovery.
- The values of recovery-oriented mental health and addictions systems are based on the recognition that each person is the agent of his or her own recovery and all services can be organized to support recovery. Person-centered services offer choice, honor each person's potential for growth, focus on a person's strengths, and attend to the overall health and wellness of a person with mental illness and/or addiction.

The authors conclude their discussion by presenting a synopsis of where the mental health and addictions fields concur on how to redesign the systems to assist people in their recovery from mental illness and/or addiction (some are extracted here). According to the authors, recovery should serve as the organizing construct for service provision and for systems improvement.

Additionally, to overcome the limitations of the current acute care model for delivering treatment services, it should be shifted to a community model, where recovery-oriented services are provided in communities, in specific environments of need, and be provided by professionals, family members, and peers. Services are structured to address the long-term and complex needs of people living with addiction and mental health issues. Moreover, to create and operate a recovery-oriented system it should include:

- Principles, e.g., multiple pathways to recovery, recovery is supported by peers, and recovery is non-linear;
- Values, e.g., person-centered services, client choice and, focus on health and wellness;
- Service strategies, e.g., treatment, post-treatment monitoring, early re-intervention, and community support; and
- Essential strategies, e.g., treatment, peer and community support, legal aid, basic support and family formation.⁵

Substance use problems and disorders are preventable and treatable chronic conditions. One aspect of providing recovery-oriented services is the shift from acute care methods to the broader adoption of chronic care strategies throughout the systems of care. Multiple articles have been published in which researchers provide a discussion of the chronic care conceptual framework and the effectiveness of addictions treatment.⁶⁻²⁷ Research in the addictions field uses the treatment system as its organizing construct. Within the recovery-oriented systems of care framework, although the treatment system is central, it is but one of multiple systems. To

illustrate this point, research suggests that the service systems that support the initiation of recovery may be different from those that sustain recovery.²⁸⁻³⁰


The life course perspective on drug use is a conceptual framework for understanding drug use trajectories. This framework classifies varying drug use trajectories, identifies critical events and factors contributing to the persistence or change in drug use, analytically orders events that occur during the lifespan, and determines contributory relationships.^{31, p.515} Hser, Longshore, and Anglin (2007), in presenting the life course perspective on drug use, discuss the evidence demonstrating the multiple service systems that drug users often come in contact with, including drug treatment, criminal justice, mental health, welfare, and primary health. They further state that the interactions with the varied social services system can “trigger turning points” for some individuals and aid in recovery.³² Turning points are changes in an existing life pathway initiated earlier in one’s life and can be positive (e.g., cohesive marriage, meaningful work) or negative (e.g., incarceration, heavy drinking or drug use). The authors’ framework incorporates the “patterns or trajectories of drug use across

individuals’ lives and the ways in which the patterns are shaped by a broader historical context and social structures.”^{33, p.517}

Babor, Stenius, and Romelsjo (2008) describe elements of a public health approach and conceptual model for the delivery of services and the service systems for people with substance use problems and disorders.³⁴ The authors primarily focus on the importance of treatment as the organizing construct of the service systems. The model includes specialized services for alcohol and drug dependence as well as medical care and social welfare services that interact with and complement specialized alcohol and drug services.³⁵ The model also includes the mediators (treatment policies) and moderators contributing to successful outcomes.

The model (Figure 2) outlines the structural resources and qualities of alcohol and drug treatment systems and includes the policy determinants and the population impact of treatment systems. The policy determinants include authoritative decisions made by governmental agencies and legislative policies, regulatory and allocative policies, treatment policies, and system qualities.

Figure 2: Conceptual Model of Population Impact of Treatment Systems



In this model:³⁶

- Treatment policies are authoritative decisions made by governmental agencies that affect the planning, financing, and monitoring of drug and alcohol services, as well as the development of a professional workforce to operate them.
- Regulatory and allocative policies are major determinants of the structural resources available to treat persons with substance use problems and disorders, including the number of facilities, the types of programs (e.g., detoxification, methadone maintenance, therapeutic communities), the settings where programs operate (e.g., hospitals, social service agencies, specialized drug and alcohol facilities) and the personnel who

work there (e.g., drug and alcohol counselors, social workers, psychiatrists, psychologists).

- Treatment policies may also affect system qualities, specifying not only where services are located, but also how they are organized and integrated. System qualities include equity (the extent to which services are equally available and accessible to all population groups), efficiency (the most appropriate mix of services), and economy (the most cost-effective services).

As demonstrated by Babor et al., individuals receive alcohol and drug services from a variety of systems, including the specialized alcohol and drug service system; the medical, psychiatric, criminal justice, and social services systems; mutual aid groups; and voluntary organizations.³⁷ Individuals

also receive informal support provided by family and friends, churches and religious organizations, workplace programs, and impaired drivers programs. There are linkages and connections between these formal and informal systems and, depending on the structural resources and system qualities, these sectors will be more or less integrated with specialized treatment and will assume a greater or lesser amount of the responsibility for managing persons with substance use problems and disorders. The authors present an interesting conceptual framework for providing addictions treatment services from a systems perspective.³⁸ This framework can contribute to the development of the broader recovery-oriented systems of care framework that focuses on providing services throughout the continuum of care and promotes and sustains individual and community recovery.

Minkoff and Cline (2004) present four characteristics of the comprehensive, continuous, integrated systems of care model for organizing services for individuals with co-occurring psychiatric and substance use problems and disorders and the eight principles of treatment for this model.³⁹ The model's four characteristics are as follows:

1. *System level of change:* The model is implemented into the entire system of care, not only for individual programs or training initiatives.
2. *Efficient use of existing resources:* The model is implemented within the context of current service resources, but emphasizes strategies between each program's requirements and environments.

3. *Incorporation of best practices:* The model is recognized by SAMHSA as a best practice for implementation for those with co-occurring psychiatric and substance disorders.
4. *Integrated treatment philosophy:* The model utilizes a common language for both the mental health and addictions fields.

Minkoff and Cline outline the eight research- and consensus-derived principles that guide the implementation of the model of care and the approach for implementing the complex multilayered system model.⁴⁰ The implementation of the model includes the following steps, which the authors detail:

1. Integrated system planning process;
2. Formal consensus on the model;
3. Formal consensus on funding the model;
4. Identification of priority populations and locus of responsibility for each;
5. Development and implementation of program standards;
6. Structures for intersystem and interprogram care coordination;
7. Development and implementation of practice guidelines;
8. Facilitation of identification, welcoming, and accessibility;
9. Implementation of continuous integrated treatment;
10. Development of basic dual diagnosis-capable competencies for all clinicians;
11. Implementation of a system-wide training plan; and
12. Development of a plan for a comprehensive program array.

There is a sizable amount of literature on recovery as the focus of recovery-oriented systems of care in the mental health field. Starting in the early 1990s, the mental health field focused on the process of recovery to guide decisions related to the mental health system. A recovery mental health model puts the locus of control and decision-making in the hands of the person who has the mental health condition.⁴¹⁻⁴³ “Recovery is pushing systems, as well as providers, to see beyond the diagnostic and categorical services, to treating the individual consumer and his/her multiple needs. The [recovery] vision ... is of an external system that reflects the internal reality of its consumers.”^{44,p.318}

Anthony (1993, 2000) describes the recovery vision and the community support system perspective that provided a framework and essential services of a recovery-oriented systems of care for mental health disorders.^{45,46} The author’s latest work outlines the relevant systems-level research that provides the foundation for his development of recovery system standard dimensions. The author delineates the essential services in recovery-oriented systems of care, including treatment, crisis intervention, case management, rehabilitation, rights protection, basic support, self-help, and wellness/prevention.⁴⁷ The standards incorporate the importance of recovery as the basis of the system and provide guidance and direction to reinforce the development of recovery-oriented systems of care. The standards are grouped by systems-level dimensions (for specific detail on the essential services and standards, please see Anthony, 2000):⁴⁸

- Design
- Evaluation
- Leadership
- Management
- Integration
- Comprehensiveness
- Consumer involvement
- Cultural relevance
- Advocacy
- Training
- Funding
- Access

Jacobson and Curtis (2000) reviewed existing literature on the conceptualizations of recovery that are integrated within recovery-oriented systems of care for the mental health field:⁴⁹

- Recovery is generally seen as a process. It does not represent a cure, but a state of being and becoming.
- The path to recovery is highly singular or unique; no two people will have identical paths or use the same benchmarks to measure their journeys.
- In contrast to the passivity of being a patient or a voiceless recipient of services, recovery is active and requires that an individual take personal responsibility for his or her own recovery, often in collaboration with friends, family, supporters, and professionals.
- A recovery orientation includes an emphasis on choice, a concept that encompasses support for autonomous action, the requirement that the individual have a range of

opportunities from which to choose and full information about those choices, and increasing personal responsibility for the consequences of choice.

- The emotional essence of recovery is hope, a promise that things can and do change, that today is not the way it will always be.
- A key theme is that of meaning, or the discovery of purpose and direction in one's life. The search for meaning is highly personal. For some people meaning may be reflected through work or social relationships. Others derive meaning from advocacy and political action. For others, the pursuit of meaning takes on strongly spiritual elements.

- Generation of hope, and
- Collaboration and partnership, e.g., consumers, professionals, and disciplines.

Finally, as discussed by Barton (1998), the three models for delivering care within a fragmented mental health system—the medical, rehabilitation, and community support system models—are responsible for the outcomes of care.⁵⁰ The recovery philosophy articulates the “process through which this occurs in partnership with the recovering consumer. From this perspective, the consumer-centered recovery philosophy is the umbrella over all models, disciplines, practices, and activities in the hospital and the community.”⁵¹ In this philosophy, system principles include:

- Empowerment of staff and consumers,
- Integration of the rehabilitation and medical model services across hospital and community settings,
- Provision of client-centered services,
- Validation of client choice,

5. Research Supporting the Principles of Recovery and Systems of Care Elements

Previous research efforts have outlined principles of effective addictions treatment. In 1999, the National Institute on Drug Abuse (NIDA) produced a research-based guide entitled *Principles of Drug Addiction Treatment*. It identified 13 principles that research has found to be associated with effective addictions treatment:⁵²

1. No single treatment is appropriate for all individuals.
2. Treatment needs to be readily available.
3. Effective treatment attends to multiple needs of the individual, not just his or her drug use.
4. An individual's treatment and services plan must be assessed continually and modified as necessary to ensure that the plan meets the person's changing needs.
5. Remaining in treatment for an adequate period of time is critical for treatment effectiveness.
6. Counseling (individual and/or group) and other behavioral therapies are critical components of effective treatment for addiction.
7. Medications are an important element of treatment for many patients, especially when combined with counseling and other behavioral therapies.
8. Addicted or drug-abusing individuals with coexisting mental disorders should have both disorders treated in an integrated way.
9. Medical detoxification is only the first stage of addictions treatment and by itself does little to change long-term drug use.
10. Treatment does not need to be voluntary to be effective.
11. Possible drug use during treatment must be monitored continuously.
12. Treatment programs should provide assessment for HIV/AIDS, hepatitis B and C, tuberculosis, and other infectious diseases, and counseling to help patients modify or change behaviors that place themselves or others at the risk of infection.
13. Recovery from drug addiction can be a long-term process and frequently requires multiple episodes of treatment.

NIDA's principles focus on the process of delivering effective treatment. The National Summit on Recovery's 12 principles of recovery provides guidelines on the process of and outcomes associated with recovery. The NIDA principles relate most closely to the following principles of recovery:

- **There are many pathways to recovery.** The pathway to recovery may include one or more episodes of psychosocial

and/or pharmacological treatment. For some, recovery involves neither treatment nor involvement with mutual aid groups. Recovery is a process of change that permits an individual to make healthy choices and improve the quality of his or her life.

- **Recovery is holistic.** Recovery is a process through which one gradually achieves greater balance of mind, body, and spirit in relation to other aspects of one's life, including family, work, and community.
- **Recovery exists on a continuum of improved health and wellness.** Recovery is not a linear process. It is based on continual growth and improved functioning.
- **Recovery involves a process of healing and self-redefinition.** Recovery is a holistic healing process in which one develops a positive and meaningful sense of identity.
- **Recovery is a reality.** It can, will, and does happen.

6. Research Supporting the Principles of Recovery

There are many pathways to recovery. Individuals are unique with specific needs, strengths, goals, health attitudes, behaviors and expectations for recovery. Pathways to recovery are highly personal, and generally involve a redefinition of identity in the face of crisis or a process of progressive change. Furthermore, pathways are often social, grounded in cultural beliefs or traditions and involve informal community resources, which provide support for sobriety. The pathway to recovery may include one or more episodes of psychosocial and/or pharmacological treatment. For some, recovery involves neither treatment nor involvement with mutual aid groups. Recovery is a process of change that permits an individual to make healthy choices and improve the quality of his or her life.

Research has shown that there are a variety of methods that assist individuals in their process of recovery.^{53,54} Some people recover naturally; others recover through treatment and/or the assistance of self-help and mutual aid groups.⁵⁵

Natural recovery involves using one's own personal resources to resolve one's addictions without the use of treatment or involvement in a mutual aid or self-help group. It is believed to be the most common recovery pathway.⁵⁶⁻⁶⁴

In studies examining the existence and success of natural recovery, alcohol was the

most studied substance, with heroin a distant second.⁶⁵ Research has shown that natural recovery is a viable and successful pathway for people with shorter and less severe substance use problems and disorders and for those with higher incomes and more stable social and occupational supports and resources.⁶⁶⁻⁷² Additionally, individuals who recover naturally typically have fewer interpersonal conflicts and rely less on avoidance coping.⁷³⁻⁷⁵

Granfield and Cloud (2001) attribute the success of natural recovery to an individual's social capital, which they define as "the benefits that accrue to an individual as a result of the network of personal contacts and associations that surround them."^{76, p.1566} The "natural communities of friends, family members, and relatives, and the social capital available through these connections, contribute significantly" to recovery success.⁷⁷

Longitudinal studies have repeatedly shown that substance use problems and disorders treatment is associated with major reductions in substance use, problems, and costs to society, and improved patient recovery.⁷⁸⁻⁸⁵ Of individuals with chronic dependence who achieved sustained recovery, the majority did so after participating in treatment—cannabis (43 percent), cocaine (61 percent), alcohol (81 percent), and heroin (92 percent).^{86,87} For some, treatment is part of the recovery process, while for others, they recover outside of the treatment system without the aid of treatment.

Participation in mutual aid groups has also been shown to be effective in supporting recovery.⁸⁸⁻⁹⁸ Laudet, Savage, and Mahmood (2002) and Scott, Dennis, and Foss (2005) found that for many individuals in long-term recovery, recovery coaches, 12-step programs, spirituality, and social and community support are integral to sustaining recovery.^{99,100} One of the most widely studied self-help/12-step groups is Alcoholics Anonymous (AA). Research has shown that patients who attend AA regularly experience better short-term alcohol-related outcomes than do patients who attend infrequently or irregularly.¹⁰¹⁻¹⁰³ Furthermore, ongoing participation in AA contributes to continued improvement of substance use disorder symptoms.¹⁰⁴⁻¹⁰⁶

Methadone is one of the most thoroughly studied pharmaceuticals in modern medicine. Methadone maintenance therapy (MMT) is one of the most widespread treatment approaches, with 179,000 individuals enrolled in MMT (out of an estimated 900,000 opiate-dependent persons in the United States) and 100,000 individuals receiving another form of treatment.¹⁰⁷ Methadone, combined with psychosocial support, leads to improved outcomes, such as decreased death rates, reduced transmission of sexually transmitted diseases, eliminated or reduced illicit opiate use, reduced criminal activity, enhanced productive behavior through employment and academic/vocational functioning, and improved global health and social functioning.¹⁰⁸

Recovery is self-directed and empowering.

While the pathway to recovery may involve one or more periods of time when activities are directed or guided

to a substantial degree by others, recovery is fundamentally a self-directed process. The person in recovery is the “agent of recovery” and has the authority to exercise choices and make decisions based on his or her recovery goals that have an impact on the process. The process of recovery leads individuals toward the highest level of autonomy of which they are capable. Through self-empowerment, individuals become optimistic about life goals.

Doty, Kasper, and Litvak (1996) and Tilly and Wiener (2001a, 2001b), reporting on the results of initiatives in several States (California, Colorado, Kansas, Maine, Michigan, Oregon, Washington, and Wisconsin), demonstrated the effectiveness of consumer-directed services.¹⁰⁹⁻¹¹¹ These evaluations show that clients who direct their own care express greater satisfaction over the services they choose. Moreover, participants in these State consumer-directed programs perceived that quality of care either improved as a result of consumer direction or at least did not suffer. An evaluation of an Arkansas consumer-directed care program found that individuals in the program were less likely than control subjects to have unmet needs, were at least as safe from adverse events and health problems, and were more likely to be satisfied with life.¹¹²

Research shows that a sense of self-efficacy is critical to successful self-management, where individuals direct and manage their own care of a variety of chronic illnesses. This is also the case in achieving improvements in health outcomes,¹¹³⁻¹¹⁶ and self-management has been

shown to be an important component of recovery from substance use.¹¹⁷

Morgenstern, Labouvie, McCrady, Kahler, and Frey (1997) found that maintaining motivation and self-efficacy for abstinence and increasing active coping post treatment were predictive of more favorable outcomes.¹¹⁸ Motivational interviewing (MI) has been shown to decrease substance use and improve outcomes.¹¹⁹ MI is based on a partnership between the provider and the individual receiving services, and it acknowledges that individual's personal responsibility and freedom of choice.¹²⁰ It replaces the concept of "resistance" with that of "ambivalence." MI allows individuals to explore their own goals and take an active role in treatment,^{121,122} thereby assisting individuals in changing their substance use behavior.¹²³ The clinician using MI assists the individual in clarifying personal goals, identifying discrepancies between the individual's current reality and his or her goals, and developing strategies to achieve those goals.^{124,125}

Recovery involves a personal recognition of the need for change and transformation.

Individuals must accept that a problem exists and be willing to take steps to address it; these steps usually involve seeking help for a substance use disorder. The process of change can involve physical, emotional, intellectual and spiritual aspects of the person's life.

Research shows that the motivation to change drug-using behavior is a major contributing factor to a successful recovery process,¹²⁶⁻¹³⁰ and the effect of motivation varies based on severity of an individual's substance use

disorder.¹³¹ Joe, Simpson, and Broome (1999) identified motivation as the best predictor of engagement and retention.¹³² The substance-using individual must accept that he or she has a problem, make a conscious choice to change, and be willing and motivated to take action to change his or her behavior.¹³³

Recovery is holistic.

Recovery is a process through which one gradually achieves greater balance of mind, body and spirit in relation to other aspects of one's life, including family, work and community.

Some literature demonstrates that the integration of the physical, emotional, and spiritual realms of an individual is influential in the quest for recovery.¹³⁴ Alternative (e.g., acupuncture) and traditional medicine approaches for substance use treatment and recovery programs promote a balance between mind, body, and spirit, and other aspects of an individual's life, and have been shown to be valuable in helping an individual achieve recovery.¹³⁵ For example, individuals who participated in a comprehensive, holistic, therapeutic community that offered many specialized services had sustained positive outcomes, including abstinence from drug and alcohol use 12 months post treatment.¹³⁶

Recovery has cultural dimensions.

Each person's recovery process is unique and impacted by cultural beliefs and traditions. A person's cultural experience often shapes the recovery path that is right for him or her.

The literature demonstrates that an individual's culture plays a vital role in his/her life and

health status and that culture must be acknowledged, addressed, and effectively utilized in recovery.¹³⁷⁻¹⁴¹ Cross-cultural studies have demonstrated the importance of delivering culturally competent care.^{142,143} Research shows that culturally competent care improves recovery and remission rates for minority populations who are at risk for relapse.¹⁴⁴⁻¹⁴⁶

Flores (2000), in his meta-analysis of research on implications of culture on clinical care, concludes that culture has significant implications on the patient-provider relationship and the delivery of efficacious care.¹⁴⁷ He further states that it is essential for providers to consider a patient's culture and linguistic issues; the failure to do so has been shown to result in inefficiencies and inequalities in patient care, such as "inaccurate histories, decreased satisfaction with care, nonadherence, poor continuity of care, less preventive screening, miscommunication,"¹⁴⁸ which leads to repercussions in care delivery.

Traditional theories of counseling and treatment are reflective of the Western cultures. It is essential that providers develop awareness, knowledge, and skills appropriate to the client's culture.^{149,150} This is particularly important when the counselor is working with individuals who do not share the counselor's racial or ethnic heritage.^{151,152} Some scholars have argued that other aspects of culture, including sexual orientation, ethnicity, religion, and heritage, are also important.^{153,154} Researchers believe that recovery rates of African-Americans, American Indians, and Asians are lower than other populations due to the failure of treatment providers and

researchers to see race as a cultural rather than a physical phenomenon.¹⁵⁵⁻¹⁵⁷

Recovery exists on a continuum of improved health and wellness.

Recovery is not a linear process. It is based on continual growth and improved functioning. It may involve relapse and other setbacks, which are a natural part of the continuum but not inevitable outcomes. Wellness is the result of improved care and balance of mind, body and spirit. It is a product of the recovery process.

The literature consistently demonstrates that for many people, substance use problems and disorders are chronic conditions involving cycles of relapse and treatment readmissions over multiple years.¹⁵⁸⁻¹⁶³ Scholars and researchers agree that recovery is a developmental and continuous process that varies from person to person.^{164,165} Individuals continuously grow and improve their functioning throughout the recovery process. Additionally, individuals benefit from lessons learned throughout the process.¹⁶⁶

Recovery emerges from hope and gratitude.

Individuals in or seeking recovery often gain hope from those who share their search for or experience of recovery. They see that people can and do overcome the obstacles that confront them and they cultivate gratitude for the opportunities that each day of recovery offers.

Several authors have written about growth beyond maintaining abstinence or

management of the problem, often describing theoretical frameworks and specific practices and techniques that help management of the problem and promote wider growth.¹⁶⁷⁻¹⁷⁵

Irving, Seidner, Burling, Pagliarini, and Robbins-Sisco (1998) found that greater hope and increased goal-oriented thinking were positively correlated to length of time abstinent, quality of life, and self-efficacy.¹⁷⁶

Additionally, listening to peers share experiences about how they dealt successfully with drug-related problems gave individuals in recovery confidence in dealing with their own situations.¹⁷⁷

In a qualitative study, Hewitt (2004) describes the posttraumatic growth that individuals experience after discontinuing alcohol and drug use.¹⁷⁸ His study expands a phenomenon reflected in the AA concept of gratitude:¹⁷⁹

recovering alcoholics often report viewing their alcoholism as a gift that brought them to a greater sense of wholeness, fulfillment, or self-actualization than they feel they would have achieved without having to confront the addiction. In his study, Hewitt found that many individuals marked a contrast between the “craziness” of their previous lives, which were devoted to drugs or alcohol, and the calmness, stability, and sanity that were more characteristic of their current post addiction lives.¹⁸⁰

The importance of having hope and believing in the possibility of a renewed sense of self and purpose is an essential component of recovery;¹⁸¹⁻¹⁸⁷ this hope must be accompanied by a desire and motivation to recover.^{188,189}

Young and Ensing (1999) found that seeking out a source of hope and inspiration helps individuals desire change and foster motivation

to embark on and/or sustain a process of change.¹⁹⁰

Recovery involves a process of healing and self-redefinition.

Recovery is a holistic healing process in which one develops a positive and meaningful sense of identity.

McMillen, Howard, Nower, and Chung (2001) identified positive by-products of the struggle with substance use problems and disorders, including changes in life priorities and increases in self-efficacy, family closeness, closeness with others, spirituality, and compassion.¹⁹¹ Many consumers view recovery as a process of discovering and fostering self-empowerment, learning self-redefinition, returning to basic functioning, and improving quality of life.¹⁹²

Recovery involves addressing discrimination and transcending shame and stigma.

Recovery is a process by which people confront and strive to overcome stigma.

Stigma and discrimination have implications for an individual’s ability to receive care and continue on the path of recovery.¹⁹³ Based on combined data from SAMHSA’s 2004 and 2005 National Survey on Drug Use and Health (NSDUH), stigma was cited as the reason for not accessing treatment by 18.5 percent of persons who needed and sought treatment but did not receive it.¹⁹⁴

Clinical practices may not be the most efficient way to reduce stigma, which is a major barrier to care, because the causes of stigma exist both within and outside of the health care

system.¹⁹⁵⁻¹⁹⁷ Furthermore, societal stigma is viewed as one of the major barriers to recovery.¹⁹⁸

While stigma is seen as a major barrier to accessing treatment, it also plays a role in affecting outcomes of treatment and access to services for individuals in recovery.¹⁹⁹⁻²⁰¹ Link, Struening, Neese-Todd, Asmussen, and Phelan (2001) found that the stigma associated with mental illness puts people more at risk for low self-esteem.²⁰² According to the *Christian Science Monitor*, experts in treatment and recovery estimate that when recovering individuals are honest about their drug histories, they will be turned down for a job 75 percent of the time.²⁰³ Additionally, a California survey found that 59 percent of employers surveyed said they would never hire anyone with a felony conviction.²⁰⁴

Recovery is supported by peers and allies.

A common denominator in the recovery process is the presence and involvement of people who contribute hope and support and suggest strategies and resources for change. Peers, as well as family members and other allies, form vital support networks for people in recovery. Providing service to others and experiencing mutual healing help create a community of support among those in recovery.

Evaluation data and research studies point to the benefits of peer-based recovery support services for consumers, individuals who provide the services, and the delivery system.²⁰⁵⁻²¹³ An individual's sustained recovery is often influenced by his or her

social interactions.²¹⁴ Peers have been shown to be integral in facilitating abstinence and preventing relapse for individuals with substance use conditions.²¹⁵⁻²²⁰ Jason, Davis, Ferrari, and Bishop (2001) and Humphreys, Huebsch, Finney, and Moos (1999) found that processes of social support mediate the transition from recovery initiation to lifelong recovery maintenance;^{221,222} furthermore, research shows that poor social supports detrimentally impact recovery and place individuals at risk for relapse.^{223,224}

Mutual aid (or "self-help") groups have been shown to play a significant role in the process of recovery.²²⁵⁻²³⁶ These peer-based support groups require no admission process or specified length of participation and are less formalized than other types of peer recovery services. The probability of stable remission rates has been shown to rise in concert with the number of recovery mutual aid groups attended in the first 3 years of recovery.²³⁷ Additionally, active and continued participation in self-help groups has been shown to improve recovery outcomes.²³⁸⁻²⁴⁶

A longitudinal study of Oxford House recovery homes suggests that receiving abstinence support, guidance, and information from recovery home members committed to the goal of long-term sobriety enhances residents' abstinence and reduces the residents' probability of relapse.^{247,248}

Recovery involves (re)joining and (re)building a life in the community.

Recovery involves a process of building or rebuilding what a person has lost or never had due to his or her condition and its consequences.

Recovery involves creating a life within the limitation imposed by that condition. Recovery is building or rebuilding healthy family, social and personal relationships. Those in recovery often achieve improvements in the quality of their life, such as obtaining education, employment and housing. They also increasingly become involved in constructive roles in the community through helping others, productive acts and other contributions.

The basic element in the process of recovery is the reclaiming of one's life in the community and the realization that one's self needs to be restored with a reawakening of old identities and establishment of new ones.²⁴⁹ Individuals entering recovery must often work to reintegrate themselves with their families and communities,²⁵⁰ while disengaging themselves from relationships, activities, and settings associated with their addictive behavior. By detaching oneself from the previous environment, an individual can find a satisfying job, non-substance-using friends, and networks of people who may be in recovery.²⁵¹⁻²⁵³ Personal resources,²⁵⁴ positive influences of family,²⁵⁵ and social and community support²⁵⁶ have been shown to be critical factors in establishing and maintaining recovery. Granfield and Cloud (2001) posited that recovery capital, which are defined as resources that support people's recovery from substance use problems and disorders, and the inclusion of a strong social network of sober friends and family members, employment, education and a range of coping skills, improves an individual's capacity to successfully recover.^{257,258}

Interventions that restructure the patient's life in the community, such as parole, methadone maintenance, and self-help groups, have also been associated with sustained abstinence.²⁵⁹ As a part of one's recovery process and to incorporate oneself into an environment that supports their recovery, many individuals make fundamental changes to their personal, professional, and social network environments.²⁶⁰⁻²⁶⁴

Recovery is a reality.

It can, will, and does happen.

Many agree that recovery is a continuous, lifelong process.²⁶⁵⁻²⁶⁷ Epidemiologic studies show that, on average, 58 percent of individuals with chronic substance dependence achieve sustained recovery.²⁶⁸⁻²⁷¹ Recovery rates for individuals with substance use problems and disorders differ by study and vary widely, from 30 percent,²⁷² 41 percent,²⁷³ 48 percent,²⁷⁴ 59 percent,²⁷⁵ 63 percent,²⁷⁶ to 72 percent.²⁷⁷

A national survey conducted on behalf of Faces and Voices of Recovery found that approximately half of individuals who self-identified as "in recovery" or "formally addicted to" alcohol and other drugs reported being in recovery more than 5 years, and 34 percent reported 10 years or more of stable recovery.²⁷⁸

7. Research Supporting the Systems of Care Elements

Person-centered

Recovery-oriented systems of care will be person-centered. Individuals will have a menu of stage-appropriate choices that fit their needs throughout the recovery process. Choices can include spiritual supports that fit with the individual's recovery needs.

A number of studies have shown that people become more committed to a course of treatment if they are allowed to choose between several alternatives rather than are forced to select a given option.²⁷⁹ Patient choice of care is important to the individual patient and improves engagement with treatment and continuing care services.²⁸⁰ Several studies have indicated that clients who were given a choice of treatment options showed greater acceptance of treatment and higher rates of recovery at follow-up.²⁸¹⁻²⁸⁶

Researchers have reported that clients who had a choice of treatment had improved treatment processes and post treatment outcomes. Individuals who were provided with the option to choose their treatment services were more likely to work harder in treatment,²⁸⁷ have more contact with their treatment program,²⁸⁸ and better adhere to program requirements than individuals who were not given a choice of treatment.²⁸⁹ Those who had treatment options were also less likely to drop out of treatment.²⁹⁰

Inclusive of family and other ally involvement

Recovery-oriented systems of care will acknowledge the important role that families and other allies can play. Family and other allies will be incorporated, when appropriate, in the recovery planning and support process. They can constitute a source of support to assist individuals in entering and maintaining recovery. Additionally, systems need to address the treatment, recovery and other support needs of families and other allies.

Research has demonstrated that involvement of concerned others can lead to improved outcomes in both alcohol²⁹¹⁻²⁹³ and drug²⁹⁴ treatment. These connections may enhance individual's self-efficacy and reduce the probability of relapse.²⁹⁵ In a review by McCrady (2004), it was shown that family and ally involvement in treatment was associated with more positive treatment outcomes in a variety of alcohol-dependent populations.²⁹⁶

Family and ally support and healthy and productive relationships nurture long-term recovery.²⁹⁷ The family has been shown to be a key determinant of an individual's commitment and ability to achieve recovery.²⁹⁸ An individual's family and other allies can be active participants, sources of strength, and resources in recovery and can

provide a central role in maintenance of recovery.²⁹⁹⁻³⁰⁴

The level of social support that an individual receives has been directly associated with engagement indicators and treatment completion.³⁰⁵ Finney, Noyes, Coutts, and Moos (1998) found that recovery-oriented support may foster greater self-efficacy and longer abstinence.³⁰⁶ The positive consequences linked to these associations may be the result of individuals acquiring effective coping strategies and greater support from other recovering individuals.³⁰⁷⁻³⁰⁹ Furthermore, it has been shown that social support, particularly through interactions with individuals in similar situations, produces positive health implications.³¹⁰

In a randomized controlled trial, it was shown that spouse involvement, regardless of the type and intensity of therapy utilized, was an effective intervention for enhancing compliance to disulfiram,³¹¹ reducing total alcohol consumption³¹²⁻³¹⁴ and maintaining treatment gains following discharge.³¹⁵ Higgins, Budney, Bickel, and Badger (1994) found that cocaine-abusing individuals participating in community reinforcement with a significant other had better outcomes than those who chose not to bring a significant other.³¹⁶

Individualized and comprehensive services across the lifespan

Recovery-oriented systems of care will be individualized, comprehensive, stage-appropriate, and flexible. Systems will adapt to the needs of individuals, rather than requiring individuals to adapt to them. They will be designed to support recovery across the lifespan. The approach to substance use disorders will

change from an acute-based model to one that manages chronic disorders over a lifetime.

Research has shown that access to and receipt of a comprehensive array of medical, psychological, and social services improves engagement, retention, and treatment outcomes.³¹⁷⁻³²⁰ For example, access to housing, employment, and legal systems has positively aided in the treatment of substance use problems and sustained abstinence.³²¹⁻³²³

Studies find that when an individual's full array of needs (e.g., food, clothing, housing, transportation, medical care, childcare, and family, psychiatric, educational, and vocational concerns) are met, short- and long-term outcomes, including retention in treatment and reduction in substance use, are improved.³²⁴⁻³²⁶ Additionally, if individuals' distinct needs are addressed, outcomes are improved.³²⁷⁻³³⁰

Similarly, individuals presenting to substance use problems and disorders treatment with comorbid psychiatric or medical conditions often have great difficulty sustaining recovery unless these conditions are fully assessed and addressed during treatment and continuing care.^{331,332}

Recovery Across the Lifespan

Due to inherent differences, issues of gender, race, and age should be considered in the treatment and recovery process.³³³⁻³³⁶ As individuals age, their health needs and social relationships evolve and their cognitive processes change.^{337,338} Treatment approaches and social support mechanisms need to be developmentally appropriate,

taking into account the age of individuals.³³⁹ Adolescent, adult, and elderly substance users differ in many ways and have unique issues and concerns that must be addressed through specific treatment and recovery planning.³⁴⁰⁻³⁴⁴ At different stages in life, an individual's drug and alcohol use often stems from different causes and requires age-appropriate treatments.³⁴⁵⁻³⁵⁰ Because of their unique developmental issues, adolescent, adult, and elderly users must be treated differently, with the variation taken into account in their treatment and recovery plans. Moreover, treatment approaches should address the nuances of each individual's experience, and their cognitive, emotional, physical, social, and moral development.^{351,352}

Research shows that the types of substances that individuals misuse varies with age. Older adults, for example, more often misuse alcohol and prescription and over-the-counter drugs, while younger substance abusers more often use illicit drugs.³⁵³⁻³⁵⁵ Adolescents are vulnerable due to their developmental stage, but multiple biological, psychological, and social changes associated with the aging process also make the elderly uniquely vulnerable to substance abuse problems.³⁵⁶⁻³⁵⁹

Natural recovery is most often seen in individuals with patterns of substance use that are moderate to mild or of short duration and is most often encountered among two age cohorts: 1) young adults whose use discontinues or is substantially reduced with maturation and assumption of adult role responsibilities, and 2) later-life adults who change behavior in response to the cumulative consequences of substance

use.^{360,361} (See also the discussion of natural recovery in Section 6.)

Systems anchored in the community

Recovery-oriented systems of care will be nested in the community for the purpose of enhancing the availability and support capacities of families, intimate social networks, community-based institutions and other people in recovery.

Research shows that social and community resources promote better recovery outcomes.³⁶² Healthy and productive environments are nurturing of recovery,^{363,364} and the presence of strong social networks during and after treatment has been linked consistently to sustained reductions in substance use after treatment.³⁶⁵⁻³⁶⁷ Strong social networks are particularly important as a counterbalance to the social pressures within high-drug-using networks and neighborhoods.³⁶⁸⁻³⁷¹

To improve individuals' long-term stability, communities must provide necessary resources, such as housing, employment, and social support.³⁷²⁻³⁷⁴ Comprehensive, easily accessible recovery support service programs located in high-need communities with staff functioning in multiple roles using culturally competent interventions have been shown to improve recovery outcomes for individuals with children.³⁷⁵ Employment and stable housing have been found to improve self-esteem and support reintegration into mainstream society and thereby to support recovery.³⁷⁶⁻³⁸¹

Community-based care systems provide opportunities for transitions between levels and types of care in a cost-effective manner and improve long-term health outcomes.^{382,383} The Team for the Assessment of Psychiatric Services (TAPS) examined the health outcomes and quality of care for individuals who were discharged into the community after extended psychiatric hospitalizations.³⁸⁴ Of 523 individuals still alive at the 5-year follow-up, 90 percent were still living in the community, and few individuals reported criminal justice system involvement or homelessness.³⁸⁵

Harrison (2001) and Knight, Simpson, and Hiller (1999) found that, among offenders reentering the community, participation in an in-prison treatment and in a post release, community-based aftercare program was associated with positive outcomes.^{386,387}

Continuity of care

Recovery-oriented systems of care will offer a continuum of care, including pretreatment, treatment, continuing care and support throughout recovery. Individuals will have a full range of stage-appropriate services from which to choose at any point in the recovery process.

Continuity of care is particularly important in treating chronic and complex diseases where several providers may be involved in the provision of care.³⁸⁸ Continuity of care is characterized by care from one doctor or team, coordinated through a common purpose and plan.³⁸⁹ Furthermore, continuity of care is significantly related to positive treatment outcomes,³⁹⁰ since continued attachment to

treatment is consistently related to better outcomes.³⁹¹

Continuing care includes services that are accessed post discharge from treatment and at a lower intensity. It provides sustained access to treatment and recovery services and promotes continued abstinence and recovery. Empirical research has demonstrated that continuing care contributes to improved treatment outcomes.^{392,393} Gruber, Fleetwood, and Herring (2001) highlighted the efficacy and the positive effects of a continuing care program designed to assist the substance-affected family.³⁹⁴ Continuing care has been shown to be effective in assisting individuals in starting and maintaining recovery.³⁹⁵ The utilization of continuing care improves long-term recovery outcomes;^{396,397} without continuing care, individuals are more likely to relapse.^{398,399}

Linkage and retention in continuing care have been shown to improve long-term abstinence from a variety of substances.⁴⁰⁰ Studies have shown that onsite medical consultation, team-based approaches, and facilitated referrals to primary care independently have a substantial positive impact on linkage to medical care and its quality for persons with mental and addictive disorders.^{401,402} For example, participation in community services has been associated with engagement in outpatient treatment and better treatment outcomes.⁴⁰³⁻

⁴¹³ Individuals who become more engaged in outpatient care in the community and self-help groups tend to experience better short-term abstinence outcomes.⁴¹⁴

Transitional services are particularly important for correctional populations.^{415,416} Substance use disorder treatment, when provided in

conjunction with credible sanctions against drug offenses, job training and placement, and advocacy services, has been shown to decrease recidivism rates and improve reentry from correctional institutions into the community for recovering drug offenders.⁴¹⁷ Completion of aftercare services by offenders has been shown to improve recovery outcomes.⁴¹⁸⁻⁴²⁰

Partnership-consultant relationships

Recovery-oriented systems of care will be patterned after a partnership-consultant model that focuses more on collaboration and less on hierarchy. Systems will be designed so that individuals feel empowered to direct their own recovery.

Research shows that supportive therapeutic and trusting relationships enhance engagement and retention.⁴²¹⁻⁴²⁴ Early therapeutic alliances appear to be a consistent predictor of engagement and retention in treatment.⁴²⁵

Providers must convey hope in their interactions with clients and develop individualized treatment or recovery plans that incorporate clients' goals and are designed to support increased patient/client autonomy.⁴²⁶

A partnership-consultant relationship is utilized to encourage patient self-management.⁴²⁷ Patient self-management requires the clinician to utilize a "collaborative care model of practice in which the patient and clinician are equal partners, with equal expertise."⁴²⁸ In the collaborative care model, the clinician brings medical expertise and "patients are experts in their own lives and in what concerns them and motivates and enables them to make changes in their lives,"⁴²⁹

thereby creating a therapeutic alliance with patients as their principal caregivers.

A therapeutic alliance between providers and patients embraces a more empathetic approach where providers enhance their clients' involvement in service delivery and recovery.⁴³⁰ Success of counseling is related to the quality of a working alliance and utilization of a patient-centered focus with therapist empathy, warmth, and genuineness.⁴³¹ A positive working alliance, reported by either the client or therapist, is a significant predictor of treatment participation and substance use behavior post treatment.^{432,433} The association of a therapeutic relationship with positive outcomes is consistently reported, and there is a positive relationship between therapeutic alliances and outcomes. Moreover, "counselors who are confrontational or use confrontational interventions consistently" have worse outcomes.⁴³⁴

Studies confirming this have been performed across settings (residential, outpatient, continuing care, and office based) and approaches (medication-assisted, adolescent and family treatment).⁴³⁵⁻⁴³⁹ Ilgen, Tiet, Finney, and Moos (2006) and Ilgen, McKellar, Moos, and Finney (2006) found that a positive therapeutic alliance counteracted the negative impact of low baseline self-efficacy and low motivation in some people.^{440,441} Carten (1996) observed that the development of positive relationships, including jointly designed service contracts, shared planning, staff encouragement, nonjudgmental attitudes, and nonpunitive responses to relapse, improved success for mothers in recovery.⁴⁴²

Strength-based

Recovery-oriented systems of care will emphasize individual strengths, assets and resiliencies.

The strengths perspective emphasizes building on the client's assets, desires, abilities, and resources to assist the client in the recovery process.⁴⁴³ Additionally, the strengths perspective demonstrates the importance and respect for the client's way of thinking and dealing with life situations. This perspective assumes that each individual has the capacity to draw from a variety of resources, skills, and motivations to focus on their strengths and create change in their lives.^{444- 447} Through examining the efficacy of combining intensive strengths-based case management services with an established program, Siegal and colleagues found that individuals who received strengths-based case management services had improved retention in treatment, lower reported drug and alcohol use, and better outcomes related to criminality and employment than those who did not receive the intervention.^{448- 450}

Culturally responsive

Recovery-oriented systems of care will be culturally sensitive, competent and responsive. There will be recognition that beliefs and customs are diverse and can impact the outcomes of recovery efforts. In addition, the cultures of those who support the recovering individual affect the recovery process.

Professional ethical guidelines and recently developed multicultural competencies for working with diverse populations suggest that

the delivery of culturally competent services is integral to the delivery of quality and effective services.⁴⁵¹ Cultural competence has become a very important focus in the health services delivery field, as demonstrated by the U.S. Department of Health and Human Services, Office of Minority Health's standards project on Culturally and Linguistically Appropriate Services (CLAS).⁴⁵²

Ignoring culture can result in many negative consequences, including missed opportunities for screening, difficulties resulting from differing responses to medication, lack of clinician knowledge about alternative and traditional remedies, diagnostic errors resulting from miscommunication, and disruptions in services.^{453- 457} Minority Americans have different health experiences than nonminorities.⁴⁵⁸ Moreover, nonminority Americans have different experiences from each other in the health care setting, even when they have similar medical conditions and insurance coverage.⁴⁵⁹⁻⁴⁶⁶

Gender- and culture-based approaches provide more effective substance abuse treatment for all individuals with substance use problems and disorders, particularly for African-Americans, Hispanics, and Asians and their families.^{467- 469}

Longshore, Grills, and Annon (1999) found that individuals in a more culturally congruent intervention were more involved in counseling sessions, more willing to self-disclose, more motivated to seek help for drug use-associated problems, and more prepared for change.⁴⁷⁰

Campbell and Alexander (2002) found that individuals treated in culturally competent substance abuse treatment practices had higher rates of specific medical and psychosocial services, such as medical exams and financial

services, but they concluded that the practices may not be uniformly effective in promoting utilization of all services.⁴⁷¹ Gender- and culture-specific care needs in relation to social structure, ethno-history, and cultural context have been shown to influence women's health and well-being as they move through recovery.⁴⁷²

Responsiveness to personal belief systems

Recovery-oriented systems of care will respect the spiritual, religious and/or secular beliefs of those they serve and provide linkages to an array of recovery options that are consistent with these beliefs.

Many researchers have documented the value of individuals' spiritual, religious, and secular beliefs in supporting recovery.⁴⁷³⁻⁴⁸¹

Religious involvement and spiritual (re)engagement appear to be correlated with and facilitate the process of recovery.⁴⁸²⁻⁴⁸⁴

Spirituality and faith, through their associations with cognitive processes, create more positive health outcomes, including optimistic life orientation, higher resilience to stress, lower levels of anxiety, and positive effective coping skills.⁴⁸⁵ Evidence shows that spirituality and faith may facilitate the process of recovery⁴⁸⁶⁻⁴⁸⁸ and promote improvements in long-term recovery.⁴⁸⁹

Spirituality, religiousness, and life meaning "enhance coping, confer hope for the future, and provide a heightened sense of control, security, and stability; they offer support and strength to resist the opportunity to use substances, all of which are very much needed to initiate and maintain recovery."

^{490,p.15} Furthermore, many individuals in recovery cite strength acquired from religion and spirituality as main factors in contributing to their long-term recovery,^{491,492} as a source of personal strength, and as a self-protection mechanism.⁴⁹³

Commitment to peer recovery support services

Recovery-oriented systems of care will include peer recovery support services. Individuals with personal experience of recovery will provide these valuable services.

Research on peer support / mutual support groups / recovering consumers as providers of alcohol and drug treatment services shows that the use of peer support is effective in helping individuals through recovery.⁴⁹⁴⁻⁴⁹⁶ Evidence shows that seeing or visualizing those similar to oneself performing activities typically increases one's belief in one's own ability to perform those activities⁴⁹⁷ and facilitates successful management of one's chronic illness.⁴⁹⁸ Moreover, peer support has been identified in the Chronic Care Model as a method to support patients in their illness self-management.^{499,500} Peer recovery support services "foster recovery in a relational, mutually-enhancing, and safe context."^{501, p.171}

Involvement in mutual aid groups provides an opportunity for individuals to participate in drug- and alcohol-free activities as role models to others, rewards their own abstinence, and helps enhance individuals' personal and social resources.⁵⁰²⁻⁵⁰⁴ Twelve-step involvement has been related to positive outcomes, including decreased substance use,⁵⁰⁵⁻⁵¹¹ enhanced psychosocial adjustment,⁵¹²⁻⁵¹⁴ and lower

health care costs.⁵¹⁵ Moreover, 12-step participation as a continuing care activity has been shown to be effective for long-term abstinence.^{516 - 519} Self-help groups help sustain recovery^{520- 523} and provide a community that strives to be drug-free with a structured mechanism for continuous abstinence or recovery.⁵²⁴

AA affiliation has been associated with self-efficacy and positive coping, which has been linked to better outcomes.^{525,526} Rational Recovery, a self-help approach started by a clinical social worker in 1986, is intended for persons with all types of addictive disorders. It adopts cognitive approaches and has been shown to be successful in engaging substance users and promoting abstinence among members.⁵²⁷

For some dual diagnosis patients, 12-step interventions have been found to be more effective in decreasing alcohol use and increasing social interactions than self-management.⁵²⁸ (For further information on self-management, please see discussion on page 16.)

Inclusion of the voices and experiences of recovering individuals and their families

The voices and experiences of people in recovery and their family members will contribute to the design and implementation of recovery-oriented systems of care. People in recovery and their family members will be included among decision-makers and have oversight responsibilities for service provision. Recovering individuals and family members will be prominently and authentically

represented on advisory councils, boards, task forces and committees at the Federal, State and local levels.

Community involvement with public health planning and implementation has been shown to be integral to improving community health.^{529,530} Additionally, with regard to improving public safety, the voices of consumers have played a large role in restructuring medical systems to better promote consumer, provider, and system safety.⁵³¹ In these movements worldwide, consumers have played key roles in defining the priorities, providing personal expertise, and reforming patient safety criteria of health care systems.⁵³² The patient safety movement is an excellent example of how partnerships among consumers and providers of care were instrumental in improving operational and systemic deficiencies.⁵³³

The rationale for seeking participation by consumers falls into three categories: 1) to improve services and decisions, 2) to gain legitimization and/or community compliance, and 3) to bring about social change with the redistribution of power or resources.⁵³⁴ Furthermore, individuals with a “particular disease become more aggressive in voicing their desire not only for more response from government and the health care system, but also for more influence over policy decisions at the macro level and treatment decisions at the micro level...[for example,] by people with disabilities and their families and those initially affected by human immunodeficiency virus/acquired immunodeficiency syndrome (HIV/AIDS).”^{535, p.216} Research shows that giving mental health consumers a significant

role in shaping services, policies, and research improves services.⁵³⁶

Some government agencies have mandated the involvement of affected populations in policy decisions through advisory groups and planning councils.⁵³⁷ Little systematic research as to the effectiveness of the involvement of consumers on decision-making has been conducted.⁵³⁸

Integrated services

Recovery-oriented systems of care will coordinate and/or integrate efforts across service systems to achieve an integrated process that responds effectively to the individual's unique constellation of strengths, desires and needs.

Integrating care has been shown to optimize recovery outcomes and improve the cost-effectiveness of delivering services.⁵³⁹⁻⁵⁴¹

Collaboration between and integration across services systems and agencies has improved the probability of individual and family recovery.⁵⁴² Research has demonstrated the efficacy and effectiveness of providing onsite primary medical care and ancillary services in the addictions treatment setting⁵⁴³⁻⁵⁴⁵ and integrating addictions services into other settings.^{546,547}

Patients in an enhanced program that used case managers who coordinated and expedited the use of medical screening, housing assistance, parenting classes, and employment services had significantly fewer physical and mental health problems, better social functioning, and less substance use at 6 months after treatment than did individuals who were not in the enhanced program.^{548,549}

Programs that are more effective with dually diagnosed individuals tend to have a combined treatment orientation; adopt a more tolerant and persuasive, rather than confrontational stance; use peer modeling in group psychotherapy; and provide continuity of care through assertive case management and after-care.⁵⁵⁰⁻⁵⁵² Additionally, individuals with co-occurring substance use problems and disorders and medical conditions have been shown to benefit from an integrated medical and substance abuse treatment program, with increased rates of abstinence, compared with individuals without co-occurring medical problems.^{553,554}

Families and children at risk often present with a complex array of needs that require the provision and utilization of multiple services.^{555,556} While effective delivery of health, child welfare, and educational services improves the lives of families who are able to access them, integration of the social services and health systems has been shown to further improve access to and provision of necessary services.⁵⁵⁷⁻⁵⁶⁰

System-wide education and training

Recovery-oriented systems of care will ensure that concepts of recovery and wellness are foundational elements of curricula, certification, licensure, accreditation and testing mechanisms. The workforce also requires continual training, at every level, to reinforce the tenets of recovery-oriented systems of care.

Educational interventions have been shown to improve physician performance and patient identification and outcomes of care.^{561,562} A

meta-analysis of disease management, which incorporated case management, found that improved disease control was associated with education of providers.⁵⁶³ Furthermore, Bukstein et al. (2005) found that continuing education is essential to providing care that is based on the latest clinical and service interventions.⁵⁶⁴

Ongoing monitoring and outreach

Recovery-oriented systems of care will provide ongoing monitoring and feedback with assertive outreach efforts to promote continual participation, re-motivation and reengagement.

Models of ongoing monitoring and early reintervention occupy a central role in the long-term management of chronic medical conditions.^{565- 568} Accumulating evidence suggests that many cases of substance use problems and disorders should be continually monitored and are best treated with “the same type and level of ongoing clinical support as other chronic illnesses.”⁵⁶⁹

The evidence shows that it is necessary to continuously evaluate and maintain connections with individuals in recovery, and by doing so, individuals at risk for relapse can reenter treatment at an earlier point of relapse.⁵⁷⁰ In general, about 50–60 percent of patients begin reusing within 6 months of treatment cessation; therefore, it is essential for ongoing monitoring and outreach to be conducted.^{571- 574} Individuals who are readmitted sooner after relapse have better short- and long-term abstinence, improved outcome measures for employment and criminality, and lower associated substance use problems.^{575,576}

Outcomes driven

Recovery-oriented systems of care will be guided by recovery-based process and outcome measures. These measures will be developed in collaboration with individuals in recovery. Outcome measures will reflect the long-term global effects of the recovery process on the individual, family and community, not just remission of biomedical symptoms. Outcomes will be measurable and include benchmarks of quality-of-life changes.

There are several entities and projects focused on measuring substance use disorders process measures and outcomes. Examples include:

- Network for the Improvement of Addiction Treatment (NIATx; <http://www.niatx.net>), a partnership between the Robert Wood Johnson Foundation’s Paths to Recovery program, the Center for Substance Abuse Treatment’s Strengthening Treatment Access and Retention (STAR) program, NIDA, and a number of independent addictions treatment organizations, and
- Washington Circle (<http://www.washingtoncircle.org>), a policy group on performance measurement for care of substance use problems and disorders

These entities have independently developed process of care and performance measures aimed to improve treatment access and retention for individuals with substance use problems and disorders.

NIATx aims to improve access to treatment through its four process measures:

1. Average time from first request to first client treatment session;
2. “No Shows,” which measures the number of patients who do not keep an appointment;
3. “Admissions,” which counts the number of unduplicated client admissions by provider; and
4. “Continuation,” which measures of the number of clients who stay engaged in treatment.

Washington Circle has and continues to develop performance measures in four domains of the process of care:

1. Prevention/Education
2. Recognition
3. Treatment
 - a. Initiation of alcohol and other plan services
 - b. Linkage of detoxification, alcohol, and other drug plans
 - c. Treatment engagement
 - d. Interventions for family members and significant others
4. Maintenance of treatment effects

Many large-scale longitudinal outcome studies have been conducted that examine various indicators of life changes, including substance use behavior, criminal behavior, education, employment, health, and social support. Examples of these data collection and outcome evaluation efforts include the Drug Abuse

Treatment Outcome Study (DATOS) conducted by NIDA, the National Treatment Improvement Evaluation Study (NTIES) by SAMHSA, the Alcohol and Drug Services Study (ADSS) by SAMHSA, and the California Outcome Study using the California Drug and Alcohol Treatment Assessment (CALDATA) by the State of California. These data sets examine the impact of treatment and behavior change processes on quality of life outcomes.

Research based

Recovery-oriented systems of care will be informed by research. Additional research on individuals in recovery, recovery venues and the processes of recovery, including cultural and spiritual aspects, is essential. Research will be supplemented by the experiences of people in recovery.

This white paper is a critical first step in examining how the research relates to the principles of recovery and systems of care elements and identifies where additional research is needed.

Adequately and flexibly financed.

Recovery-oriented systems of care will be adequately financed to permit access to a full continuum of services, ranging from detoxification and treatment to continuing care and recovery support. In addition, funding will be sufficiently flexible to permit unbundling of services, enabling the establishment of a customized array of services that can evolve over time in support of an individual's recovery.

Pooling of funding may reduce problems associated with fragmented funding and separate service systems. Pooled funding also may improve service coordination within and between different organizations and networks and may work to expand access to and provision of services.⁵⁷⁷ Additionally, States and service facilities can utilize creative financing mechanisms to provide and reimburse for recovery-oriented services. For example, Michigan's State agencies have organized multiple funding sources to provide integrated community mental health and primary medical care, and have improved access to these services.⁵⁷⁸

Voucher programs provide people seeking drug and alcohol treatment and recovery support services with a funding mechanism to pay for a range of unbundled community-based services. Many States, through the

Access to Recovery grant program, have implemented voucher programs to improve Access to Recovery support services and promote patient choice in service provision.

8. Research Supporting the Implementation of Recovery-Oriented Services and Systems of Care

While many States (e.g., Connecticut, Georgia, Massachusetts, Michigan, New Jersey, New York, and Oregon) and cities (e.g., Philadelphia) are in the process of reforming their systems to implement recovery-oriented systems of care, there are few publications in peer-reviewed journals that describe the effectiveness of this systems change for individuals with substance use problems and disorders. However, there are several published articles in peer-reviewed journals that discuss the implementation and delivery of recovery-oriented services in the mental health system. This section discusses the available peer-reviewed literature focusing on the delivery of recovery-oriented services and the implementation of recovery-oriented systems of care in the addictions and mental health fields.

In 2000, the State of Connecticut embarked on transforming its system of publicly funded behavioral health care into a system that is recovery-oriented and culturally responsive. In 2002, Connecticut was the first State that incorporated recovery as the overarching aim of its publicly funded system of care.⁵⁷⁹ The initiative targeted the statewide system of care as a whole instead of supplementing the existing system with recovery-oriented elements.⁵⁸⁰ The Connecticut initiative involved several interrelated steps occurring over several years:

1. Developing core values and principles based on the input of people in recovery;
2. Establishing a conceptual framework based on this vision of recovery;
3. Building workforce competencies and skills through training, education, and consultation;
4. Changing programs and service structures;
5. Aligning fiscal and administrative policies in support of recovery; and
6. Monitoring, evaluating, and adjusting the efforts.

Two peer-reviewed articles describe the implementation efforts of the Connecticut system. A 2005 article in *Psychiatric Rehabilitation Journal* outlines the results of a recovery self-assessment implemented in Connecticut, which gauged perceptions of the degree to which programs implement recovery-oriented practices.⁵⁸¹ In this article, O'Connell et al. provide a statewide assessment of recovery-oriented practices in agencies. Results indicated that mental health professionals, persons in recovery, and family members "generally agreed that their agencies were providing services that are consistent with a recovery orientation."⁵⁸² A 2007 article in *Psychiatric Rehabilitation Journal* provides evaluative details on the implementation of the early stages of the initiative.⁵⁸³ Two themes are discussed:

1. Recovery does not refer to any specific service, intervention, or support (which can be more or less recovery oriented) but to what people with psychiatric disabilities or addiction *do* themselves in order to manage their illness and/or get their lives back.
2. Recovery cannot be an “add-on” to existing services, supports, or systems. The focus of transformation needs to be on changing and realigning current policies, practices, procedures, services, and supports to be oriented toward, and effective in, promoting recovery.

In 2004, Philadelphia created the city’s Department of Behavioral Health and Mental Retardation Services, which combined the Community Behavioral Health Office, the Office of Mental Health, and the Coordinating Office for Drug and Alcohol Abuse Programs into an integrated behavioral health care system.⁵⁸⁴ The systems transformation in Philadelphia included numerous stakeholders in the process, including recovering people and their

families, in developing a vision for the Philadelphia system. This vision and mission of Philadelphia was to create an integrated system that “promotes long-term recovery, resiliency, self-determination, and a meaningful life in the community,” which shifted the values of the system from the “interventions of professional experts to the experience and needs of recovering individuals and their families.”^{585, p.36}

Relationships between service practitioners, service consumers, the department, and local services providers shifted to partnerships based on mutual respect and collaborations. Additionally, “recovery representation” was emphasized at all levels of the system to “affirm that recovery is a living reality in the City of Philadelphia.”⁵⁸⁶ Practices of delivery services were changed to improve care delivery and focus on recovery. As discussed by White (2007) in his description of the recovery revolution in Philadelphia, in 10 service areas changes to the system occurred. These areas are the following: engagement, assessment, retention, role of client, service relationship, clinical care, service dose/duration, service delivery sites, post treatment checkups and support, and attitude toward readmission.⁵⁸⁷

**Modification of Philadelphia Department of Behavioral Health and Mental Retardation Services Service System and Clinical Practice To Be More Recovery Oriented
(White, 2007, pp. 36-27)**

1. *Engagement*: Greater focus on early identification via outreach and community education; emphasis on removing personal and environmental obstacles to recovery; shift in responsibility for motivation to change from the client to service provider; loosening of admission criteria; renewed focus on the quality of the service relationship.
2. *Assessment*: Greater use of global and strength-based assessment instruments and interview protocol; shift from assessment as an intake activity to assessment as a continuing activity focused on the developmental stages of recovery.
3. *Retention*: Increased focus on service retention and decreasing premature service disengagement; use of outreach workers, recovery coaches, and advocates to reduce rates of client disengagement and administrative discharge.
4. *Role of Client*: Shift toward philosophy of choice rather than prescription of pathways and styles of recovery; greater client authority and decision-making within the service relationship; emphasis on empowering clients to self-manage their own recoveries.
5. *Service Relationship*: Service relationships are less hierarchical with counselor serving more as ongoing recovery consultant than professional expert; more a stance of "How can I help you?" than "This is what you must do."
6. *Clinical Care*: Greater accountability for delivery of services that are evidence-based, gender-sensitive, culturally competent, and trauma informed; greater integration of professional counseling and peer-based recovery support services; considerable emphasis on understanding and modifying each client's recovery environment; use of formal recovery circles (recovery support network development).
7. *Service Dose/Duration*: Dose and duration of total services will increase while number and duration of acute care episodes will decline; emphasis shifts from crisis stabilization to ongoing recovery coaching; great value placed in continuity of contact in a primary recovery support relationship over time.
8. *Service Delivery Sites*: Emphasis on transfer of learning from institutional to natural environments; greater emphasis on home-based and neighborhood-based service delivery; greater use of community organization skills to build or help revitalize indigenous recovery supports where they are absent or weak.
9. *Post-treatment Checkups and Support*: Emphasis on recovery resource development (e.g., supporting alumni groups and expansion/diversification of local recovery support groups); assertive linkage to communities of recovery; face-to-face, telephone-based, or Internet-based post-treatment monitoring and support; stage-appropriate recovery education; and, when needed, early re-intervention.
10. *Attitude toward Re-admission*: Returning clients are welcomed (not shamed); emphasis on transmitting principles and strategies of chronic disease management; focus on enhancement of recovery maintenance skills rather than recycling through standard programs focused on recovery initiation; emphasis on enhancing peer-based recovery supports and minimizing need for high-intensity professional services.

Finally, Philadelphia worked with multiple constituents to plan and implement changes in funding and regulatory policies, which are critical to effectively implement and sustain the behavioral health system, and to focus its regulatory and policy reform on recovery. Philadelphia has focused on “providing regulatory relief (reducing duplicative and excessive regulatory requirements), generating more recovery-focused regulatory standards, shifting the focus of program monitoring from one of policing to one of consultation and support, generating new Request for Proposals for recovery-focused service initiatives, and exploring models for long-term funding of recovery support services.”⁵⁸⁸

The State of Massachusetts, through its Departments of Mental Health and Public Health, designed a collaborative model to provide a comprehensive integrated service system for persons with co-occurring substance use problems and disorders and serious mental illness for the private and public sectors. Barreira, Espey, Fishbein, Moran, and Flannery (2000) describe the design phase of Massachusetts’ integrated system and provide lessons learned and outcomes for developing the system, but the implementation of the initiative has not yet been published.⁵⁸⁹ Specifically, the authors discuss utilizing a collaborative framework, which includes involving the key stakeholders, to foster change and buy-in for the system. The collaborative and consensus-building process builds on evidence-based practices and best practices, builds on organizational strengths, and is sensitive to barriers to change (e.g., differing philosophies, regulatory processes, clinical

and administrative traditions and policies, resistance to change) and collaboration. Massachusetts also discovered that “providing a voice for stakeholders who are parties in designing change” and “developing consensus on a framework of care that all groups endorse” so funding resources are not seen as being taken from one system to another were critical for the collaborative process to succeed and the integrated framework to be implemented.⁵⁹⁰

Jacobson and Curtis, in a 2000 *Psychiatric Rehabilitation Journal* article, outline how the concept of recovery is being implemented in the policies and practices of several State mental health systems and review specific strategies that States, including Wisconsin, Ohio, Vermont, and Nebraska, are using to implement recovery principles into their mental health systems.⁵⁹¹ Most of the State systems reviewed begin their systems’ transformation with the development of vision statements in consensus workgroups and task forces, renaming programs, and applying strategies for operationalizing recovery-oriented services. Both programmatic and administrative strategies are being adopted to implement recovery-oriented services; they include education, consumer and family involvement, support for consumer-operated services, emphasis on relapse prevention and management, incorporation of crisis planning and advance directives, implementing stigma reduction initiatives, innovations in contracting and financing mechanisms, definition and measurement of outcomes, and reviewing and revising key policies. In this article, while Jacobson and Curtis do not specifically review activities that occur across different service systems, they discuss the importance of

involving a variety of stakeholders in the systems-change effort and of recovery education initiatives and the destigmatization that is occurring across workers in different disciplinary backgrounds, consumers, family members, and administrators. They further state that the concept of recovery must look beyond the service provider and the mental health system.

In a 2004 article that outlines the complementary nature of evidence-based practices and recovery in a service system, Solomon and Stanhope describe the profound changes required at the systems level for the implementation of evidence-based practices and recovery-oriented services.⁵⁹² The authors trace the integration of a recovery orientation into the Ohio Department of Mental Health. In 1993, Ohio started the process to transform its system. The first step of the transformation was to have a dialog with providers, consumers, and family members to explore the mental health recovery process and prepare a report outlining the stages of the process. The stages were integrated into

a framework for implementing recovery-oriented practices, which included clinical care, peer and family support, facilitation of employment, empowerment, stigma reduction, community involvement, access to resources, and education. By offering grants to localities, Ohio helped localities transform their mental health systems through establishing recovery centers and recovery management plans within agencies. “The Ohio recovery model is an example of tailoring a recovery vision to the specific needs of consumers and implementing change through financial incentives.”^{593, p.318} Other States, including California, Massachusetts, New York, and Washington, have written recovery principles within their State managed care contracts. These principles “require that organizations contract with providers who pursue recovery-oriented services, including consumer-operated services, and that consumers have an advisory role on managed care organization boards.”^{594, p.319}

9. Conclusion

This white paper examines the research that supports the principles of recovery and systems of care elements as defined by the National Summit on Recovery. The author identified findings in more than 375 studies that supported the framework, principles, elements, and implementation of recovery-oriented services and systems. This document is intended to serve as a starting point for further examination of recovery research. Additionally, it provides States, communities, and organizations that are developing and implementing recovery-oriented services and systems with evidence that supports services and systems improvements based on recovery-oriented approaches.

While many of the principles and systems elements are easily supported by existing literature in the addictions field, research supporting others was more difficult to find. In some circumstances, they were supported by literature outside of addictions research, primarily through the mental health and public health research fields. What follows is a brief synopsis of the research that was found to support the principles of recovery and systems of care elements and a listing of the fields from which the information was derived.

Extensive research has been conducted in the addictions field to support the following principles and systems elements:

- There are many pathways to recovery;

- Recovery exists on a continuum of improved health and wellness;
- Recovery is supported by peers and allies;
- Recovery is a reality;
- Inclusive of family and other ally involvement;
- Individualized and comprehensive services across the lifespan;
- Continuing care part of the continuity of care element;
- Partnership-consultant relationships;
- Responsiveness to personal belief systems;
- Commitment to peer recovery support services;
- Integrated services; and
- Ongoing monitoring and outreach.

The following principles and systems elements were supported by a modest amount of research from the addictions field:

- Recovery is self-directed and empowering;
- Recovery involves a personal recognition of the need for change and transformation;
- Recovery emerges from hope and gratitude;

- Recovery involves addressing discrimination and transcending shame and stigma;
- Recovery involves (re)joining and (re)building a life in the community;
- Systems anchored in the community;
- Strength-based; and
- Outcomes driven.

The following principles and systems elements were supported by limited addictions research, but were grounded in literature from the general public health and mental health fields:

- Recovery has cultural dimensions;
- Person-centered;
- Continuity of care;
- Culturally responsive;
- Inclusion of the voices and experiences of recovering individuals and their families; and
- System-wide education and training.

The following principles were supported by a minimal amount of research in the addictions, mental health, and public health research fields:

- Recovery is holistic;
- Recovery involves a process of healing and self-redefinition; and
- Adequately and flexibly financed.

In relation to the process- and outcomes-driven systems element, it should be noted that large-scale, longitudinal studies have been conducted that measure quality-of-life outcomes for individuals with substance use problems and disorders. However, this systems element points to the importance of examining long-term global effects of the recovery process on the individual, family, and community, while also studying the process of care and implementing continuous quality improvement mechanisms to improve treatment access and retention. Additionally, this element suggests that process and outcome measures should be developed in collaboration with individuals in recovery. Although the addictions research community includes some researchers in recovery, this element suggests broader inclusion of the voices of individuals in recovery in developing outcome measures.

Finally, this paper identifies areas where there are modest amounts of addictions research related to recovery. Limited literature on the conceptual framework of recovery-oriented systems of care exists, but even less research is available on the implementation and outcomes of recovery-oriented services and systems for those with substance use problems and disorders. By providing this baseline assessment and the associated gaps in the research, future research agendas can be better informed.

References

- 1 Center for Substance Abuse Treatment. (2005). *National Summit on Recovery: Conference Report*. DHHS Publication No. (SMA) 07-4276. Rockville, MD: Substance Abuse and Mental Health Services Administration.
- 2 Anthony, W.A. (2000). A recovery-oriented service system: setting some system level standards. *Psychiatric Rehabilitation Journal*, 24(2), 159–168.
- 3 O’Connell, M., Tondora, J., Croog, G., Evans, A., & Davidson, L. (2005). From rhetoric to routine: Assessing perceptions of recovery-oriented practices in a state mental health and addiction system. *Psychiatric Rehabilitation Journal*, 28(4), 378–386.
- 4 Gagne, C., White, W., Anthony, W.A. (2007) Recovery: A common vision for the fields of mental health and addictions. *Psychiatric Rehabilitation Journal*, 31(1): 32-37.
- 5 Ibid.
- 6 Friedmann, P.D., Hendrickson, J.C., Gerstein, D.R., & Zhang, Z. (2004). The effect of matching comprehensive services to patients’ needs on drug use improvement in addiction treatment. *Addiction*, 99(8), 962–972.
- 7 Simpson, D.D., Joe, G.W., Greener, J.M., & Rowan-Szal, G.A. (2000). Modeling year 1 outcomes with treatment process and post-treatment social influences. *Substance Use and Misuse*, 35, 1911–1930.
- 8 Joe, G.W., Broome, K.M., Rowan-Szal, G.A., & Simpson, D.D. (2002). Measuring patient attributes and engagement in treatment. *Journal of Substance Abuse Treatment*, 22(4), 183–196
- 9 Hser, Y.I., Evans, E., Huang, D., & Anglin, D.M. (2004). Relationship between drug treatment services, retention, and outcomes. *Psychiatric Services*, 55(7), 767–774.
- 10 French, M.T., McGeary, K.A., Chitwood, D.D., & McCoy, C.B. (2000). Chronic illicit drug use, health services utilization and the cost of medical care. *Social Science & Medicine*, 50, 1703–1713.
- 11 French, M.T., Salome, H.J., Sindelar, J.L., & McLellan, A.T. (2002). Benefit-cost analysis of addiction treatment: Methodological guidelines and empirical application using the DATCAP and ASI. *Health Services Research*, 37, 433–455.
- 12 Hser, Y.I., Hoffman, V., Grella, C.E., Anglin, M.D. (2001). A 33-year follow-up of narcotics addicts. *Archives of General Psychiatry*, 58, 503–508.
- 13 Etheridge, R.M, Craddock, S.G., Dunteman, G.H., & Hubbard, R.L. (1995). Treatment services in two national studies of community-based drug abuse treatment programs. *Journal of Substance Abuse*, 7(1), 9–26.
- 14 Salome, H.J., French, M.T., Miller, M., & McLellan, A.T. (2003). Estimating the client costs of addiction treatment: First findings from the client drug abuse treatment cost analysis program (Client DATCAP). *Drug and Alcohol Dependence*, 71, 195–206.
- 15 Simpson, D.D., & Sells, S.B. (1974). Patterns of multiple drug abuse: 1969-1971. *International Journal of Addictions*, 9, 301–314.
- 16 Simpson, D.D., Joe, G.W., & Rowan-Szal, G.A. (1997). Drug abuse treatment retention and process effects on follow-up outcomes. *Drug and Alcohol Dependence*, 47, 227–235.
- 17 Simpson, D.D., Joe, G.W., Fletcher, B.W., Hubbard, R.L., & Anglin, M.D. (1999). A national evaluation of treatment outcomes for cocaine dependence. *Archives of General Psychiatry*, 56, 507–514.
- 18 Simpson, D.D. (2004). A conceptual framework for drug treatment process and outcomes. *Journal of Substance Abuse Treatment*, 27, 99–121.
- 19 McLellan, A.T., Lewis, D.C., O’Brien, C.P., & Kleber, H.D. (2000). Drug dependence, a chronic medical illness: Implications for treatment, insurance, and outcomes evaluation. *Journal of the American Medical Association*, 284(13), 1689–1695.
- 20 McLellan, A.T., Woody, G.E., Metzger, D., McKay, J., Alterman, A.I., & O’Brien, C.P. (1996). Evaluating the effectiveness of addiction treatments: Reasonable expectations, appropriate comparisons. *Milbank Quarterly*, 74(1), 51–85.
- 21 Simpson, D.D., & Brown, B.S. (Eds.) (1999). Treatment process and outcome studies from DATOS [Special issue]. *Drug and Alcohol Dependence*, 57.
- 22 Simpson, D.D., & Curry, S.J. (Eds.) (1997). Drug Abuse Treatment Outcome Study (DATOS) [Special issue]. *Psychology of Addictive Behaviors*, 11.

- 23 Simpson, D.D., & Sells, S.B. (1982). Effectiveness of treatment for drug abuse: An overview of the DARP research program. *Advances in Alcohol and Substance Abuse*, 2, 7–29.
- 24 Lamb, S., Greenlick, M.R., & McCarty, D. (1998). *Bridging the gap between practice and research: Forging partnerships with community-based drug and alcohol treatment*. Washington, DC: National Academy Press.
- 25 Leshner, A.I. (1997). Drug abuse and addiction treatment research. The next generation. *Archives of General Psychiatry*, 54, 691–694.
- 26 McLellan, A.T., Lewis, D.C., O'Brien, C.P., & Kleber, H.D. (2000). Drug dependence, a chronic medical illness: Implications for treatment, insurance, and outcomes evaluation. *Journal of the American Medical Association*, 284, 1689–1695.
- 27 Haggerty, J.L., Reid, R.J., Freeman, G.K., Starfield, B.H., Adair, C.E., & McKendry, R. (2003). Continuity of care: A multidisciplinary review. *British Medical Journal*, 327, 1219–1221.
- 28 Humphreys, K., Moos, R.H., & Finney, J.W. (1995). Two pathways out of drinking problems without professional treatment. *Addictive Behaviors*, 20(4), 427–441.
- 29 White, W.L., Boyle, M., & Loveland, D. (2002). Alcoholism/addiction as a chronic disease: From rhetoric to clinical reality. *Alcoholism Treatment Quarterly*, 20(3/4), 107–130.
- 30 Laudet, A.B., Savage, R., & Mahmood, D. (2002). Pathways to long-term recovery: A preliminary investigation. *Journal of Psychoactive Drugs*, 34(3), 305–311.
- 31 Hser, Y., Longshore, D., & Anglin, M.D. (2007). The life course perspective on drug use: A conceptual framework for understanding drug use trajectories. *Evaluation Review*, 31(6), 515–547.
- 32 Ibid.
- 33 Ibid, p. 517.
- 34 Babor, T.F., Stenius, K., & Romelsjo, A. (2008). Alcohol and drug treatment systems in public health perspective: Mediators and moderators of population effects. *International Journal of Methods in Psychiatric Research*, 17(S1), S50–S59.
- 35 Ibid.
- 36 Ibid.
- 37 Ibid.
- 38 Ibid.
- 39 Minkoff, K., & Cline, C.A. (2004). Changing the world: The design and implementation of comprehensive continuous integrated systems of care for individuals with co-occurring disorders. *Psychiatric Clinics of North America*, 27(4), 727–743.
- 40 Minkoff, K., & Cline, C.A. (2004). Changing the world: The design and implementation of comprehensive continuous integrated systems of care for individuals with co-occurring disorders. *Psychiatric Clinics of North America*, 27(4), 727–743.
- 41 Frese, F.F., Stanley, J., Kress, K., & Vogel-Scibilia, S. (2001). Integrating evidence-based practices and the recovery model. *Psychiatric Services*, 52(11), 1462–1468.
- 42 Jacobson, N., & Greenley, D. (2001). What is recovery? A conceptual model and explication. *Psychiatric Services*, 52, 482–485.
- 43 Jacobson, N., & Curtis, L. (2000). Recovery as policy in mental health services: Strategies emerging from the states. *Psychiatric Rehabilitation Journal*, 23(4), 333–341.
- 44 Solomon, P., & Stanhope, V. (2004). Recovery: Expanding the vision of evidence-based practice. *Brief Treatment and Crisis Intervention*, 4(4), 311–321.
- 45 Anthony, W.A. (1993). Recovery from mental illness: The guiding vision of the mental health service system in the 1990s. *Psychiatric Rehabilitation Journal*, 16(4), 11–23.
- 46 Anthony W.A. (2000). A recovery-oriented service system: Setting some system level standards. *Psychiatric Rehabilitation Journal*, 24(2), 159–168.
- 47 Anthony W.A. (2000). A recovery-oriented service system: Setting some system level standards. *Psychiatric Rehabilitation Journal*, 24(2), 159–168.
- 48 Ibid.
- 49 Jacobson, N., & Curtis, L. (2000).
- 50 Barton, R. (1998). The rehabilitation-recovery paradigm: A statement of philosophy for a public mental health system. *Psychiatric Rehabilitation Skills*. 2(2), 171–187.

- 51 Ibid, p. 177.
- 52 National Institute on Drug Abuse. (1999; reprinted 2000). *Principles of drug addiction treatment: A research based guide* (NIH Publication No. 00-4180). Bethesda, MD: National Institutes of Health.
- 53 Laudet, A.B., Savage, R., & Mahmood, D. (2002).
- 54 Venner, K.L., Matzger, H., Forcehimes, A.A., Moos, R.H., Feldstein, S.W., Willenbring, M.L., et al. (2006). Course of recovery from alcoholism. *Alcoholism, Clinical and Experimental Research, 30*, 1079–1090.
- 55 Laudet, A.B., Savage, R., & Mahmood, D. (2002).
- 56 Havassy, B.E., Hall, S.M., & Wasserman, D.A. (1991). Social support and relapse: Commonalities among alcoholics, opiate users, and cigarette smokers. *Addictive Behaviors, 16*, 235–246.
- 57 Sobell, L.C., Sobell, M.B., Toneatto, T., & Leo, G.I. (1993). Severely dependent alcohol abusers may be vulnerable to alcohol cues in television programs. *Journal of Studies on Alcohol, 54*, 85–91.
- 58 Fillmore, K.M., Grant, M., Hartka, E., Johnstone, B.M., Sawyer, S.M., Speigman, R., et al. (1988). Collaborative longitudinal research on alcohol problems. *British Journal of Addictions, 83*, 441–444.
- 59 Sobell, L.C., Sobell, M.B., Toneatto, T., & Leo, G.I. (1993). What triggers the resolution of alcohol problems without treatment. *Alcoholism, Clinical and Experimental Research, 17*, 217–224.
- 60 Cunningham, J.A., Sobell, L.C., Sobell, M.B., & Kapur, G. (1995). Resolution from alcohol problems with and without treatment: Reasons for change. *Journal of Substance Abuse, 7*(3), 365–72.
- 61 Cunningham, J.A. (1999). Untreated remissions from drug use: The predominant pathway. *Addictive Behaviors, 24*, 267–270.
- 62 Cunningham, J. A. (1999). Resolving alcohol-related problems with and without treatment: The effects of different problem criteria. *Journal of Studies on Alcohol, 60*, 463–466.
- 63 Sobell, L.C., Cunningham, J.A., & Sobell, M.B. (1996). Recovery from alcohol problems with and without treatment: Prevalence in two population surveys. *American Journal of Public Health, 86*, 966–972.
- 64 Sobell, L.C., Ellingstad, T.P., & Sobell, M.B. (2000). Natural recovery from alcohol and drug problems: Methodological review of the research with suggestions for future directions. *Addiction, 95*, 749–764.
- 65 Ibid.
- 66 Sobell, L.C., Sobell, M.B., Toneatto, T., et al (1993).
- 67 Sobell, L.C., Cunningham, J.A., & Sobell, M.B. (1996).
- 68 Larimer, M.E., & Kilmer, J.R. (2000). Natural history. In G. Zernig, A. Saria, M. Kurz, & S.S. O'Malley (Eds.), *Handbook of alcoholism* (pp. 13–28). Boca Raton, FL: CRC Press.
- 69 Latkin, C.A., Knowlton, A.R., Hoover, D., & Mandell, W. (1999). Drug network characteristics as a predictor of cessation of drug use among adult injection drug users: A prospective study. *The American Journal of Drug and Alcohol Abuse, 25*(3), 463–473.
- 70 Weisner, C., Matzger, H., & Kaskutas, L.A. (2003). How important is treatment? One-year outcomes of treated and untreated alcohol-dependent individuals. *Addiction, 98*, 901–911.
- 71 Bischof, G., Rumpf, H.J., Hapke, U., Meyer, C., & John, U. (2001). Factors influencing remission from alcohol dependence without formal help in a representative population sample. *Addiction, 96*, 1327–1336.
- 72 Russell, M., Peirce, R.S., Chan, A.W.K., Wieczorek, W.F., Moscato, B.S., & Nochajski, T.H. (2001). Natural recovery in a community-based sample of alcoholics: Study design and descriptive data. *Substance Use and Misuse, 36*(11), 1417–1441.
- 73 Weisner, C., Matzger, H., & Kaskutas, L.A. (2003).
- 74 Bischof, G., Rumpf, H.J., Hapke, U., Meyer, C., & John, U. (2001).
- 75 Russell, M., Peirce, R.S., Chan, A.W.K., Wieczorek, W.F., Moscato, B.S., & Nochajski, T.H. (2001).
- 76 Granfield, R., & Cloud, W. (2001). Social context and “natural recovery”: The role of social capital in the resolution of drug-associated problems. *Substance Use and Misuse, 36*, 1543–1570.
- 77 Ibid, p. 1566.
- 78 Etheridge, R.M, Craddock, S.G., Dunteman, G.H., & Hubbard, R.L. (1995).
- 79 Simpson, D.D., & Sells, S.B. (1974).
- 80 Simpson, D.D., Joe, G.W., & Rowan-Szal, G.A. (1997).
- 81 Simpson, D.D., Joe, G.W., Fletcher, B.W., Hubbard, R.L., & Anglin, M.D. (1999).
- 82 French, M.T., McGeary, K.A., Chitwood, D.D., & McCoy, C.B. (2000). Chronic illicit drug use, health services utilization and the cost of medical care. *Social Science & Medicine, 50*, 1703–1713.

- 83 French, M.T., Salome, H.J., Sindelar, J.L., & McLellan, A.T. (2002). Benefit-cost analysis of addiction treatment: Methodological guidelines and empirical application using the DATCAP and ASI. *Health Services Research, 37*, 433-455.
- 84 Hser, Y.I., Hoffman, V., Grella, C.E., & Anglin, M.D. (2001). A 33-year follow-up of narcotics addicts. *Archives of General Psychiatry, 58*, 503-508.
- 85 Salome, H.J., French, M.T., Miller, M., & McLellan, A.T. (2003). Estimating the client costs of addiction treatment: First findings from the client drug abuse treatment cost analysis program (Client DATCAP). *Drug and Alcohol Dependence, 71*, 195-206.
- 86 Cunningham, J.A., Koski-Jannes, A., & Toneatto, T. (1999). Why do people stop their drug use? Results from a general population sample. *Contemporary Drug Problems, 26*, 695-710.
- 87 Cunningham, J.A., Lin, E., Ross, H.E., & Walsh, G.W. (2000). Factors associated with untreated remissions from alcohol abuse or dependence. *Addictive Behaviors, 25*, 317-321.
- 88 Laudet, A.B., Savage, R., & Mahmood, D. (2002).
- 89 Miller, W.R., & Longabaugh, R. (2002). Summary and conclusions. In T.F. Babor & F.K. Del Boca (Eds.), *Treatment matching in alcoholism*. Cambridge, UK: Cambridge University Press.
- 90 Ouimette, P.C., Moos, R.H., & Finney, J.W. (1998). Influence of outpatient treatment and 12-step group involvement on one-year substance abuse treatment outcomes. *Journal of Studies on Alcohol, 59*, 513-522.
- 91 Scott, C.K., Dennis, M.L., & Foss, M.A. (2005). Utilizing recovery management checkups to shorten the cycle of relapse, treatment reentry, and recovery. *Drug and Alcohol Dependence, 78*, 325-338.
- 92 Florentine, R. (1999). After drug treatment: Are 12-step programs effective in maintaining abstinence? *American Journal of Drug and Alcohol Abuse, 25*, 93-116.
- 93 Miller, N.S., Ninonuevo, F.G., Klamen, D.L., Hoffmann, N.G., & Smith, D.E. (1997). Integration of treatment and posttreatment variables in predicting results of abstinence-based outpatient treatment after one year. *Journal of Psychoactive Drugs, 29*, 239-248.
- 94 McKellar, J., Kelly, J., Harris, A., & Moos, R. (2006). Pretreatment and during treatment risk factors for dropout among patients with substance use disorders. *Addictive Behaviors, 31*(3), 450-60. Epub 2005 Jun 23.
- 95 Tonigan, J.S., Miller, W.R., & Connors, G.J. (2001). The search for meaning in life as a predictor of alcoholism treatment outcome. In R. Longabaugh & P.W. Wirtz (Eds.), *Project MATCH Monograph Series: Vol. 8. Project MATCH hypotheses: Results and causal chain analysis* (pp. 154-165). Bethesda, MD: National Institute on Alcohol Abuse and Alcoholism.
- 96 Connors, G.J., Tonigan, J.S., & Miller, W.R. (2001). A longitudinal model of intake symptomatology, AA participation and outcome: Retrospective study of the project MATCH outpatient and aftercare samples. *Journal of Studies on Alcohol, 62*, 817-825.
- 97 Johnson, N.P., & Chappel, J.N. (1994). Using AA and other 12-step programs more effectively. *Journal of Substance Abuse Treatment, 11*(2), 137-142.
- 98 Khantzian, E.J., & Mack, J.E. (1994). How AA works and why it's important for clinicians to understand. *Journal of Substance Abuse Treatment, 11*(2), 77-92.
- 99 Laudet, A.B., Savage, R., & Mahmood, D. (2002).
- 100 Scott, C.K., Dennis, M.L., & Foss, M.A. (2005).
- 101 Tonigan, J.S., Miller, W.R., & Connors, G.J. (2001).
- 102 Connors, G.J., Tonigan, J.S., & Miller, W.R. (2001). A longitudinal model of intake symptomatology, AA participation and outcome: Retrospective study of the project MATCH outpatient and aftercare samples. *Journal of Studies on Alcohol, 62*, 817-825.
- 103 Johnson, N.P., & Chappel, J.N. (1994). Using AA and other 12-step programs more effectively. *Journal of Substance Abuse Treatment, 11*(2), 137-142.
- 104 Tonigan, J.S., Miller, W.R., & Connors, G.J. (2001).
- 105 Connors, G.J., Tonigan, J.S., & Miller, W.R. (2001). A longitudinal model of intake symptomatology, AA participation and outcome: Retrospective study of the project MATCH outpatient and aftercare samples. *Journal of Studies on Alcohol, 62*, 817-825.
- 106 Johnson, N.P., & Chappel, J.N. (1994). Using AA and other 12-step programs more effectively. *Journal of Substance Abuse Treatment, 11*(2), 137-142.
- 107 Kreek, M.J. & Vocci, F.J. (2002). History and current status of opioid maintenance treatments: Blending conference session. *Journal of Substance Abuse Treatment, 23*(2), 93-105.

- 108 National Consensus Development Panel on Effective Medical Treatment of Opiate Addiction. (1998). Effective medical treatment of opiate addiction. *JAMA*, *280*, 1936–1943.
- 109 Doty, P., Kasper, J., & Litvak, S. (1996). Consumer-directed models of personal care: Lessons from Medicaid. *Milbank Quarterly*, *74*(3): 377–409.
- 110 Tilly, J., & Wiener, J. (2001a). Consumer-directed home and community services programs in eight states: Policy issues for older people and government. *Journal of Aging and Social Policy*, *12*(4),1–2.
- 111 Tilly, J., & Wiener, J. (2001b). *Consumer-directed home and community services: Policy issues*. Washington, DC: The Urban Institute.
- 112 Foster, L., Brown, R., Phillips, B., Schore, J., Lepidus, R., & Carlson, B. (2003). Improving the quality of Medicaid personal assistance through consumer direction. *Health Affairs Web Exclusive* [serial online], Mar 26. Retrieved from <http://www.healthaffairs.org/WebExclusives/FosterWebExcl032603.htm>
- 113 Lorig, K.R., & Holman, H. (2003). Self-management education: History, definition, outcomes, and mechanisms. *Annals of Behavioral Medicine*, *26*, 1–7.
- 114 Lorig, K.R., Ritter, P., Stewart, A.L., Sobel, D.S., Brown, B.W., Bandura, A., et al. (2001). Chronic disease self-management program: 2-year health status and health care utilization outcomes. *Medical Care*, *39*, 1217–1223.
- 115 Shoor, S., & Lorig, K.R. (2002). Self-care and the doctor-patient relationship. *Medical Care*, *40*, 1140–1144.
- 116 Institute of Medicine. (2006). *Improving the quality of health care for mental and substance use conditions*. Washington, DC: National Academy of Sciences.
- 117 Samet, J.H., Rollnick, S., & Barnes, H. (1996). Beyond CAGE: A brief clinical approach after detection of substance abuse. *Archives of Internal Medicine*, *156*, 2287–2293.
- 118 Morgenstern, J., Labouvie, E., McCrady, B.S., Kahler, C.W., & Frey, R.M. (1997). Affiliation with Alcoholics Anonymous after treatment: A study of its therapeutic effects and mechanisms of action. *Journal of Consulting and Clinical Psychology*, *65*, 768–777.
- 119 Miller, W.R. (1996). Motivational interviewing: Research, practice, and puzzles. *Addictive Behaviors*, *21*, 835–842.
- 120 Swanson, A.J., Pantalon, M.V., & Cohen, K.R. (1999). Motivational interviewing and treatment adherence among psychiatric and dually diagnosed patients. *Journal of Nervous and Mental Disease*, *187*, 630–635.
- 121 Miller, W.R., & Rollnick, S. (1991). *Motivational interviewing: Preparing people to change addictive behavior*. New York: Guilford Press.
- 122 Miller, W.R., & Rollnick, S. (2002). *Motivational interviewing: Preparing people for change* (2nd ed.). New York: Guilford Press.
- 123 Rusch, N., & Corrigan, P.W. (2002). Motivational interviewing to improve insight and treatment adherence in schizophrenia. *Psychiatric Rehabilitation Journal*, *26*, 23–32.
- 124 Miller, W.R., & Rollnick, S. (1991).
- 125 Miller, W.R., & Rollnick, S. (2002).
- 126 Venner, K.L., Matzger, H., Forcehimes, A.A., et al. (2006).
- 127 Maddux, J.F., & Desmond, D.P. (1986). Relapse and recovery in substance abuse careers. *NIDA Research Monograph*, *72*, 49–71.
- 128 Miller, W.R. (1998). Researching the spiritual dimensions of alcohol and other drug problems. *Addiction*, *93*, 979–990.
- 129 Flynn, P.M., Joe, G.W., Broome, K.M., Simpson, D.D., & Brown, B.S. (2003). Recovery from opioid addiction in DATOS. *Journal of Substance Abuse Treatment*, *25*, 177–186.
- 130 Drake, R.E., Wallach, M.A., & McGovern, M.P. (2005). Future directions in preventing relapse to substance abuse among clients with severe mental illnesses. *Psychiatric Services*, *56*, 1297–1302.
- 131 Carpenter, K.M., Miele, G.M., & Hasin, D.S. (2002). Does motivation to change mediate the effect of DSM-IV substance use disorders on treatment utilization and substance use? *Addictive Behaviors*, *27*(2), 207–225.
- 132 Joe, G.W., Simpson, D.D., & Broome, K.M. (1999). Retention and patient engagement models for different treatment modalities in DATOS. *Drug and Alcohol Dependence*, *57*, 113–125.
- 133 Laudet, A. B., Morgen, K., & White, W. L. (2006). The role of social supports, spirituality, religiousness, life meaning and affiliation with 12-step fellowships in quality of life satisfaction among individuals in recovery from alcohol and drug problems. *Alcoholism Treatment Quarterly*, *24*, 33–73.

- 134 Lowery, C.T. (1998). American Indian perspectives on addiction and recovery. *Health and Social Work, 23*, 127–135.
- 135 Winkelman, M. (2003). Complementary therapy for addiction: “Drumming out drugs.” *American Journal of Public Health, 93*, 647–651.
- 136 Carroll, J.F., & McGinley, J.J. (2000). An agency follow-up outcome study of graduates from four inner-city therapeutic community programs. *Journal of Substance Abuse Treatment, 18*, 103–118.
- 137 Wingo, L.K. (2001). Substance abuse in African American women. *Journal of Cultural Diversity, 8*, 21–25.
- 138 Peterson, S., Berkowitz, G., Cart, C.U., & Brindis, C. (2002). Native American women in alcohol and substance abuse treatment. *Journal of Health Care for the Poor and Underserved, 13*, 360–378.
- 139 Hyun, M.S., Kools, S., & Kim, S.A. (2003). A model of recovery from substance abuse and dependence for Korean adolescents. *Journal of Child and Adolescent Psychiatric Nursing, 16*(1), 25–34.
- 140 Bowser, B.P., & Bilal, R. (2001). Drug treatment effectiveness: African-American culture in recovery. *Journal of Psychoactive Drugs, 33*, 391–402.
- 141 Grunbaum, J.A., Tortolero, S., Weller, N., & Gingiss, P. (2000). Cultural, social, and intrapersonal factors associated with substance use among alternative high school students. *Addictive Behaviors, 25*, 145–151.
- 142 Harwood A. (1981). *Ethnicity and medical care*. Cambridge, MA: Harvard University Press.
- 143 Kleinman, A., Eisenberg, L., & Good, B. (1978). Culture, illness, and care: Clinical lessons from anthropologic and crosscultural research. *Annals of Internal Medicine, 88*, 251–258.
- 144 Scott, C.K., Dennis, M.L., & Foss, M.A. (2005).
- 145 Wingo, L.K. (2001).
- 146 Bowser, B.P., & Bilal, R. (2001).
- 147 Flores, G. (2000). Culture and the patient-physician relationship: Achieving cultural competency in health care. *The Journal of Pediatrics, 136*, 14–23.
- 148 Ibid.
- 149 Sue, D.W., Arrendondo, P., & McDavis, R.J. (1992). Multicultural counseling competencies and standards: A call to the profession. *Journal of Counseling and Development, 70*, 477–486.
- 150 Sue, D.W., Bernier, Y., Durran, A., Feinberg, L., Pedersen, P.B., Smith, E.J., et al. (1982). Position paper: Cross-cultural counseling competencies. *Counseling Psychologist, 10*, 45–52.
- 151 Sue, D.W., Arrendondo, P., & McDavis, R.J. (1992).
- 152 Sue, D.W., Bernier, Y., Durran, A., Feinberg, L., et al. (1982).
- 153 Croteau, J.M., Evans, N.J., Lance, T.S., & Talbot, D.M. (2002). A qualitative study of the interplay between privilege and oppression. *Journal of Multicultural Counseling and Development, 30*(4): 239–258.
- 154 Matthews, C.R., Lorah, P., & Fenton, J. (2006). Treatment experiences of gays and lesbians in recovery from addiction: A qualitative inquiry. *Journal of Mental Health Counseling, 28*(2), 111–132.
- 155 Wingo, L.K. (2001).
- 156 Hyun, M.S., Kools, S., & Kim, S.A. (2003).
- 157 Bowser, B.P., & Bilal, R. (2001).
- 158 Anglin, M.D., Hser, Y.I., Grella, C.E., Longshore, D., & Prendergast, M.L. (2001). Drug treatment careers: Conceptual overview and clinical, research, and policy applications. In F. Tims, C. Leukefeld, & J. Platt (Eds.), *Relapse and recovery in addictions* (pp. 18–39). New Haven, CT: Yale University Press.
- 159 Dennis, M.L., Scott, C.K., & Hristova, L. (2002). The duration and correlates of substance abuse treatment careers among people entering publicly funded treatment in Chicago. *Drug and Alcohol Dependence, 66*(Suppl. 2), 44.
- 160 Hser, Y.I., Anglin, M.D., Grella, C., Longshore, D., & Prendergast, M.L. (1997). Drug treatment careers. A conceptual framework and existing research findings. *Journal of Substance Abuse Treatment, 14*, 543–558.
- 161 Lamb, S., Greenlick, M.R., & McCarty, D. (1998). *Bridging the gap between practice and research: Forging partnerships with community-based drug and alcohol treatment*. Washington, DC: National Academy Press.
- 162 Leshner, A.I. (1997). Drug abuse and addiction treatment research. The next generation. *Archives of General Psychiatry, 54*, 691–694.
- 163 McLellan, A.T., Lewis, D.C., O'Brien, C.P., & Kleber, H.D. (2000). Drug dependence, a chronic medical illness: Implications for treatment, insurance, and outcomes evaluation. *JAMA, 284*, 1689–1695.
- 164 Drake, R.E., Wallach, M.A., & McGovern, M.P. (2005).

- 165 Rawson, R.A., Obert, J.L., McCann, M.J., & Ling, W. (1991). Psychological approaches for the treatment of cocaine dependence: A neurobehavioral approach. *Journal of Addictive Disorders, 11*, 97–119.
- 166 Drake, R.E., Wallach, M.A., & McGovern, M.P. (2005).
- 167 Irving, L.M., Seidner, A.L., Burling, T.A., Pagliarini, R., & Robbins-Sisco, D. (1998). Hope and recovery from substance dependence in homeless veterans. *Journal of Social and Clinical Psychology, 17*(4), 389–406.
- 168 Magura, S., Knight, E.L., Vogel, H.S., Mahmood, D., Laudet, A.B., & Rosenblum, A. (2003). Mediators of effectiveness in dual-focus self-help groups. *American Journal of Drug and Alcohol Abuse, 29*, 301–322.
- 169 Bewley, A.R. (1993). Addiction and meta-recovery: Wellness beyond the limits of Alcoholics Anonymous. *Alcoholism Treatment Quarterly, 10*(1/2), 1–22.
- 170 Whitfield, C.L. (1984). Stress management and spirituality during recovery: A transpersonal approach. Part II: Being. *Alcoholism Treatment Quarterly, 1*(2).
- 171 Whitfield, C.L. (1984). Stress management and spirituality during recovery: A transpersonal approach. Part III: Transforming. *Alcoholism Treatment Quarterly, 1*(4).
- 172 Grof, S. (1987). Spirituality, addiction, and western science. *ReVision, 10*(2), 18.
- 173 Small, J. (1987). Spiritual emergence and addiction: A transpersonal approach to alcoholism and drug abuse counseling. *ReVision, 10*(2), 23–36.
- 174 Sparks, T. (1987). Transpersonal treatment of addictions: Radical return to roots. *ReVision, 10*(2), 49–64.
- 175 McMillen, C., Howard, M.O., Nower, L., & Chung, S. (2001). Positive by-products of the struggle with chemical dependency. *Journal of Substance Abuse Treatment, 20*, 69–79.
- 176 Irving, L.M., Seidner, A.L., Burling, T.A., et al. (1998).
- 177 Magura, S., Knight, E.L., Vogel, H.S., et al. (2003).
- 178 Hewitt, A. (2004). *Post-traumatic growth in substance misuse*. Annegatan, Finland: Nordic Counsel for Alcohol and Drug Research.
- 179 Whitfield, C.L. (1984). Stress management and spirituality during recovery: A transpersonal approach. Part I: Becoming. *Alcoholism Treatment Quarterly, 1*(1).
- 180 Hewitt, A. (2004).
- 181 Davidson, L., Stayner, D.A., Nickou, C., Styron, T.H., Rowe, M., & Chinman, M.L. (2001). Simply to be let in: Inclusion as a basis for recovery. *Psychiatric Rehabilitation Journal, 24*, 375–388.
- 182 Deegan, P.E. (1996). Recovery as a journey of the heart. *Psychiatric Rehabilitation Journal, 19*, 91–97.
- 183 Fisher, D. (1994). Health care reform based on an empowerment model of recovery by people with psychiatric disabilities. *Hospital and Community Psychology, 45*(9), 913–915.
- 184 Jacobson, N., & Curtis, L. (2000). Recovery as policy in mental health services: Strategies emerging from the states. *Psychiatric Rehabilitation Journal, 23*(4), 333–341.
- 185 Jacobson, N., & Greenley, D. (2001). What is recovery? A conceptual model and explication. *Psychiatric Services, 52*(4), 482–485.
- 186 Mead, S., & Copeland, M.E. (2000). What recovery means to us: Consumers' perspectives. *Community Mental Health Journal, 36*(3), 315–328.
- 187 Smith, M.K. (2000). Recovery from a severe psychiatric disability: Findings of a qualitative study. *Psychiatric Rehabilitation Journal, 24*(2), 149–159.
- 188 Young, S.L., & Ensing, D.S. (1999). Exploring recovery from the perspective of people with psychiatric disabilities. *Psychiatric Rehabilitation Journal, 22*, 219–231.
- 189 Substance Abuse and Mental Health Services Administration. (2006). *Results from the 2005 National Survey on Drug Use and Health: National findings* (NSDUH Series H-30, DHHS Publication No. SMA 06-4194). Rockville, MD: Office of Applied Studies.
- 190 Young, S.L., & Ensing, D.S. (1999).
- 191 McMillen, C., Howard, M.O., Nower, L., et al. (2001).
- 192 Young, S.L., & Ensing, D.S. (1999).
- 193 Institute of Medicine. (2006).
- 194 Substance Abuse and Mental Health Services Administration. (2006).
- 195 Office of the Surgeon General. (2001). *Mental Health: Culture, race, and ethnicity: A supplement to mental health: A report of the Surgeon General*. Rockville, MD: U.S. Department of Health and Human Services, Public Health Services.

- 196 National Institute of Mental Health. (1998). *Bridging science and service: A report by the National Advisory Mental Health Council's Treatment and Services Workgroup*. Bethesda, MD: National Institute of Mental Health.
- 197 Wells, K., Miranda, J., Bruce, M.L., Alegria, M., & Wallerstein, N. (2004). Bridging community intervention and mental health services research. *American Journal of Psychiatry*, 161, 955–963.
- 198 Perlick, D.A. (2001). Special section on stigma as a barrier to recovery: Introduction. *Psychiatric Services*, 52, 1613–1614.
- 199 Link, B.G., Struening, E.L., Neese-Todd, S., Asmussen, S., & Phelan, J.C. (2001). Stigma as a barrier to recovery: The consequences of stigma for the self-esteem of people with mental illnesses. *Psychiatric Services*, 52(12), 1621–1626.
- 200 Marks, A. (2002). Jobs elude former drug addicts. *Christian Science Monitor*. Retrieved from <http://www.csmonitor.com/2002/0604/p02s02-ussc.html>
- 201 Curley, B. (2002). Discrimination against people in recovery rampant, advocates say. *Join Together Online*. Retrieved from <http://www.jointogether.org/sa/news/features/reader/0,1854,553416,00.html>
- 202 Link, B.G., Struening, E.L., Neese-Todd, S., et al. (2001).
- 203 Marks, A. (2002).
- 204 Curley, B. (2002).
- 205 Humphreys, K., Wing, S., McCarty, D., Chappel, J., Gallant, L., Haberle, B., et al. (2004). Self-help organizations for alcohol and drug problems: Toward evidence-based practice and policy. *Journal of Substance Abuse Treatment*, 26, 151–158.
- 206 Brown, S.A., D'Amico, E.J., McCarthy, D.M., & Tapert, S.F. (2001). Four-year outcomes from adolescent alcohol and drug treatment. *Journal of Studies on Alcohol*, 62, 381–388.
- 207 Brown, S.A., Gleghorn, A., Schuckit, M.A., Myers, M.G., & Mott, M.A. (1996). Conduct disorder among adolescent alcohol and drug abusers. *Journal of Studies on Alcohol*, 57, 314–324.
- 208 Catalano, R.F., Hawkins, J.D., Wells, E.A., Miller, J., & Brewer, D. (1991). Evaluation of the effectiveness of adolescent drug abuse treatment, assessment of risks for relapse, and promising approaches for relapse prevention. *International Journal of the Addictions*, 25, 1085–1140.
- 209 Richter, S.S., Brown, S.A., & Mott, M.A. (1991). The impact of social support and self-esteem on adolescent substance abuse treatment outcome. *Journal of Substance Abuse*, 3, 371–385.
- 210 Shoemaker, R.H., & Sherry, P. (1991). Posttreatment factors influencing outcome of adolescent chemical dependency treatment. *Journal of Adolescent Chemical Dependency*, 2(1), 89–106.
- 211 Solomon, P. (2004). Peer support/peer provided services underlying processes, benefits, and critical ingredients. *Psychiatric Rehabilitation Journal*, 27, 392–401.
- 212 Solomon, P., & Draine, J. (2001). The state of knowledge of the effectiveness of consumer provided services. *Psychiatric Rehabilitation Journal*, 25, 20–27.
- 213 Christensen, A., & Jacobsen, N.S. (1994). Who (or what) can do psychotherapy: The status and challenge of nonprofessional therapies. *Psychological Science*, 5, 8–14.
- 214 Venner, K.L., Matzger, H., Forcehimes, A.A., et al. (2006).
- 215 Humphreys, K., Wing, S., McCarty, D., et al. (2004).
- 216 Brown, S.A., D'Amico, E.J., McCarthy, D.M., et al. (2001).
- 217 Brown, S.A., Gleghorn, A., Schuckit, M.A., et al. (1996).
- 218 Catalano, R.F., Hawkins, J.D., Wells, E.A., et al. (1991).
- 219 Richter, S.S., Brown, S.A., & Mott, M.A. (1991).
- 220 Shoemaker, R.H., & Sherry, P. (1991).
- 221 Jason, L.A., Davis, M.I., Ferrari, J.R., & Bishop, P.D. (2001). Oxford House: A review of research and implications for substance abuse recovery and community research. *Journal of Drug Education*, 31, 1–27.
- 222 Humphreys, K., Huebsch, P.D., Finney, J.W., & Moos, R.H. (1999). A comparative evaluation of substance abuse treatment: Substance abuse treatment can enhance the effectiveness of self-help groups. *Alcoholism, Clinical and Experimental Research*, 23, 558–563.
- 223 Venner, K.L., Matzger, H., Forcehimes, A.A., et al. (2006).
- 224 Wingo, L.K. (2001).
- 225 Florentine, R. (1999).

- 226 Morgenstern, J., Labouvie, E., McCrady, et al. (1997).
- 227 Jason, L.A., Davis, M.I., Ferrari, J.R., et al. (2001).
- 228 Humphreys, K., Huebsch, P.D., Finney, J.W., et al. (1999).
- 229 Jason, L.A., Davis, M.I., & Ferrari, J.R. (2007). The need for substance abuse after-care: Longitudinal analysis of Oxford House. *Addictive Behaviors*, 32, 803–818.
- 230 Mankowski, E.S., Humphreys, K., & Moos, R.H. (2001). Individual and contextual predictors of involvement in twelve-step self-help groups after substance abuse treatment. *American Journal of Community Psychology*, 29, 537–563.
- 231 Timko, C., Moos, R.H., Finney, J.W., & Moos, B.S. (1994). Outcome of treatment for alcohol abuse and involvement in Alcoholics Anonymous among previously untreated problem drinkers. *Journal of Mental Health Administration*, 21(2), 145–160.
- 232 Fiorentine, R., & Hillhouse, M.P. (2000). Exploring the additive effects of drug misuse treatment and twelve-step involvement: Does Twelve-Step ideology matter? *Substance Use and Misuse*, 35, 367–397.
- 233 Emrick, C.D., Tonigan, J.S., Montgomery, H., & Little, L. (1993). Alcoholics Anonymous: What is currently known. In B.S. McCrady & W.R. Miller (Eds.), *Research on Alcoholics Anonymous* (pp. 41–76). Brunswick, NJ: Rutgers Center of Alcohol Studies.
- 234 Tucker, J.A., Vuchinich, R.E., & Gladsjo, J.A. (1994). Environmental events surrounding natural recovery from alcohol-related problems. *Journal of Studies on Alcohol*, 55, 401–411.
- 235 McKay, J.R., McLellan, A.T., Alterman, A.I., Cacciola, J.S., Rutherford, M.J., & O'Brien, C.P. (1998). Predictors of participation in aftercare sessions and self-help groups following completion of intensive outpatient treatment for substance abuse. *Journal of Studies on Alcohol*, 59, 152–162.
- 236 Fiorentine, R., & Hillhouse, M. (2000). Drug treatment and 12-step participation: The additive effects of integrated recovery activities. *Journal of Substance Abuse Treatment*, 18(1), 65–74.
- 237 Hoffmann, N., Harrison, P., & Belille, C. (1983). Alcoholics Anonymous after treatment: Attendance and abstinence. *International Journal of the Addictions*, 18(3), 311–318.
- 238 Fiorentine, R. (1999).
- 239 Fiorentine, R., & Hillhouse, M.P. (2000).
- 240 Emrick, C.D., Tonigan, J.S., Montgomery, H., et al. (1993).
- 241 Hoffmann, N., Harrison, P., & Belille, C. (1983).
- 242 Cohen, C., Humphreys, K., & Moos, R.H. (1997). Social and community resources and long-term recovery from treated and untreated alcoholism. *Journal of Studies on Alcohol*, 58(3), 231–238.
- 243 Emrick, C.D. (1987). Alcoholics Anonymous: Affiliation processes and effectiveness as treatment. *Alcoholism, Clinical and Experimental Research*, 11, 416–423.
- 244 Kingree, J.B. (1995). Understanding gender differences in psychosocial functioning and treatment retention. *American Journal of Drug and Alcohol Abuse*, 21(2), 267–281.
- 245 McKay, J.R., Alterman, A.I., McLellan, A.T., & Snider, E.C. (1994). Treatment goals, continuity of care, and outcome in a day hospital substance abuse rehabilitation program. *American Journal of Psychiatry*, 151, 254–259.
- 246 Timko, C., & Sempel, J.M. (2004). Intensity of acute services, self-help attendance and one-year outcomes among dual diagnosis patients. *Journal of Studies on Alcohol*, 65, 274–282.
- 247 Jason, L.A., Davis, M.I., Ferrari, J.R., et al. (2001).
- 248 Jason, L.A., Davis, M.I., & Ferrari, J.R. (2007).
- 249 Biernacki, P. (1986). *Pathways from heroin addiction: Recovery without treatment*. Philadelphia: Temple University Press.
- 250 Gruber, K.J., Fleetwood, T.W., & Herring, M.W. (2001). In-home continuing care services for substance-affected families: The bridges program. *Social Work*, 46, 267–277.
- 251 Miller, W.R. (1998).
- 252 Landry, M.J. (1994). *Understanding drugs of abuse: The processes of addiction, treatment, and recovery*. Arlington, VA: American Psychiatric Publishing.
- 253 Davis, K.E., & O'Neill, S.J. (2005). A focus group analysis of relapse prevention strategies for persons with substance use and mental disorders. *Psychiatric Services*, 56, 1288–1291.

- 254 Moos, R. H. (1994). Why do some people recover from alcohol dependence, whereas others continue to drink and become worse over time? *Addiction*, *89*, 31–34.
- 255 Miller, W.R., & Rollnick, S. (2002).
- 256 Cohen, C., Humphreys, K., & Moos, R.H. (1997).
- 257 Granfield, R., & Cloud, W. (2001).
- 258 Ibid.
- 259 Vaillant, G.E. (1988). What can long-term follow-up teach us about relapse and prevention of relapse in addiction? *British Journal of Addictions*, *83*, 1147–1157.
- 260 Miller, W.R. (1996).
- 261 Landry, M.J. (1994).
- 262 Davis, K.E., & O'Neill, S.J. (2005).
- 263 Moos, R. H. (1994).
- 264 Vaillant, G.E. (1988).
- 265 Miller, W.R., & Rollnick, S. (2002).
- 266 Flynn, P.M., Joe, G.W., Broome, K.M., et al. (2003).
- 267 Margolis, R., Kilpatrick, A., & Mooney, B. (2000). A retrospective look at long-term adolescent recovery: Clinicians talk to researchers. *Journal of Psychoactive Drugs*, *32*, 117–125.
- 268 Margolis, R., Kilpatrick, A., & Mooney, B. (2000).
- 269 Kessler, R.C. (1994). The National Comorbidity Survey of the United States. *International Review of Psychiatry*, *6*, 365–376.
- 270 Dawson, D.A. (1996). Correlates of past-year status among treated and untreated persons with former alcohol dependence: United States, 1992. *Alcoholism, Clinical and Experimental Research*, *20*, 771–779.
- 271 Robins, L.N., & Regier, D.A. (1991). *Psychiatric disorders in America: The Epidemiologic Catchment Area Study*. New York: Free Press.
- 272 Schutte, K.K., Nichols, K.A., Brennan, P.L., & Moos, R.H. (2003). A ten-year follow-up of older problem drinkers: Risk of relapse and implications of successfully sustained remission. *Journal of Studies on Alcohol*, *64*(3), 367–374.
- 273 Ojesjo, L. (1981). Long-term outcome in alcohol abuse alcoholism among males in the Lundby general population, Sweden. *British Journal of Addictions*, *76*, 391–400.
- 274 Dawson, D.A., Grant, B.F., Stinson, F.S., Chou, P.S., Huang, B., & Ruan, W.J. (2005). Recovery from DSM-IV alcohol dependence: United States, 2001–2002. *Addiction*, *100*, 281–292.
- 275 Vaillant, G.E. (2003). A 60-year follow-up of alcoholic men. *Addiction*, *98*, 1043–1051.
- 276 Helzer, J.E., Burnam, A., & McEvoy, L.T. (1991). Alcohol abuse and dependence. In L.N. Robins & D.A. Regier (Eds.), *Psychiatric disorders in America: The Epidemiologic Catchment Area Study* (pp. 81–115). New York: Free Press.
- 277 Dawson, D.A. (1996).
- 278 Faces and Voices of Recovery. (2001). *The road to recovery: A landmark national study on the public perceptions of alcoholism and barriers to treatment*. San Francisco: Peter D. Hart Research Associates / The Recovery Institute.
- 279 Brigham, T. (1979). Some effects of choice on academic performance. In L.C. Perlmutter & R.A. Monty (Eds.), *Choice and perceived control*. Hillsdale, NJ: Lawrence Erlbaum.
- 280 Laugharne, R., & Priebe, S. (2006). Trust, choice and power in mental health: A literature review. *Social Psychiatry and Psychiatric Epidemiology*, *41*(11), 843–852.
- 281 Langer, E., & Rodin, J. (1976). The effects of choice and enhanced personal responsibility for the aged: A field experiment in an institutional setting. *Journal of Personality and Social Psychology*, *34*, 191–198.
- 282 Calsyn, R.J., Winter, J.P., & Morse, G.A. (2000). Do consumers who have a choice of treatment have better outcomes? *Community Mental Health Journal*, *36*, 149–160.
- 283 Thompson, C.E., & Wankel, L.M. (1980). The effects of perceived activity choice upon frequency of exercise behavior. *Journal of Applied Psychology*, *10*, 436–443.
- 284 Rokke, P.D., Tomhave, J.A., & Jovic, Z. (1999). The role of client choice and target selection in self-management therapy for depression in older adults. *Psychology of the Aging*, *14*(1), 155–169.

- 285 Kissin, B., Platz, A., & Su, W.H. (1971). Selective factors in treatment choice and outcome in alcoholics. In N.K. Mello and J.K. Mendelson (Eds.), *Recent advances in studies of alcoholism* (NIMH Publication No. [HSM] 71-9045, pp. 781-802). Washington, DC: U.S. Government Printing Office.
- 286 Miller, W.R. (1985). Motivation for treatment: A review with special emphasis on alcoholism. *Psychological Bulletin*, *98*, 84-107.
- 287 Langer, E., & Rodin, J. (1976).
- 288 Calsyn, R.J., Winter, J.P., & Morse, G.A. (2000).
- 289 Thompson, C.E., & Wankel, L.M. (1980).
- 290 Rokke, P.D., Tomhave, J.A., & Jovic, Z. (1999).
- 291 McCrady, B.S., Noel, N.E., Abrams, D.B., Stout, R.L., Nelson, H.F., & Hay, W.M. (1986). Comparative effectiveness of three types of spouse involvement in outpatient behavioral alcoholism treatment. *Journal of Studies on Alcohol*, *47*(6), 459-467.
- 292 O'Farrell, T.J., Cutter, H.S.G., & Floyd, F.J. (1985). Evaluating behavioral marital therapy for male alcoholics: Effects on marital adjustment and communication from before to after therapy. *Behavior Therapy*, *16*, 147-167.
- 293 Noel, N.E., McCrady, B.S., Stout, R.L., & Fisher-Nelson, H. (1987). Predictors of attrition from an outpatient alcoholism treatment program for couples. *Journal of Studies on Alcohol*, *48*(3), 229-235.
- 294 Higgins, S.T., Budney, A.J., Bickel, W.K., & Badger, G.J. (1994). Participation of significant others in outpatient behavioral treatment predicts greater cocaine abstinence. *The American Journal of Drug and Alcohol Abuse*, *20*, 47-56.
- 295 Jason, L.A., Davis, M.I., & Ferrari, J.R. (2007).
- 296 McCrady, B.S. (2004). To have but one true friend: Implications for practice of research on alcohol use disorders and social networks. *Psychology of Addictive Behaviors*, *18*, 113-121.
- 297 Drake, R.E., Wallach, M.A., & McGovern, M.P. (2005).
- 298 Gruber, K.J., & Fleetwood, T.W. (2004). In-home continuing care services for substance use affected families. *Substance Use and Misuse*, *39*, 1379-1403.
- 299 Ibid.
- 300 Brown, S., & Lewis, V. (1998). *The Alcoholic Family in Recovery. A Developmental Model*. New York: The Guilford Press.
- 301 Carten, A.J. (1996). Mothers in recovery: Rebuilding families in the aftermath of addiction. *Social Work*, *41*, 214-223.
- 302 Donovan, D.M. (1998). Continuing care: Promoting the maintenance of change. In: W.R. Miller & N. Heather (Eds.), *Treating addictive behaviors* (2nd ed.). New York: Plenum Press.
- 303 Higgins, C., Duxbury, L., & Lee, C. (1994). Impact of life-cycle stage and gender on the ability to balance work and family responsibilities. *Family Relations*, *43*(2), 144-150.
- 304 O'Farrell, T.J., Choquette, K.A., Cutter, H.S., Brown, E.D., & McCourt, W.F. (1993). Behavioral marital therapy with and without additional couples relapse prevention sessions for alcoholics and their wives. *Journal of Studies on Alcohol*, *54*, 652-666.
- 305 Simpson, D. & Joe, G. (2003). A longitudinal evaluation of treatment engagement and recovery stages. *Journal of Substance Abuse Treatment*, *27*, (2), 89-97.
- 306 Finney, J.W., Noyes, C.A., Coutts, A.I., & Moos, R.H. (1998). Evaluating substance abuse treatment process models: Changes on proximal outcome variables during 12-step and cognitive-behavioral treatment. *Journal of Studies on Alcohol*, *59*, 371-380.
- 307 Granfield, R., & Cloud, W. (2001).
- 308 Laudet, A. B., Morgen, K., & White, W. L. (2006).
- 309 Weisner, C., Delucchi, K., Matzger, H., & Schmidt, L. (2003). The role of community services and informal support on five-year drinking trajectories of alcohol dependent and problem drinkers. *Journal of Studies on Alcohol*, *64*, 862-873.
- 310 Berkman, L.F. (2000). Social support, social networks, social cohesion and health. *Social Work and Health Care*, *31*, 3-14.
- 311 Azrin, N.H., Sisson, R.W., Meyers, R., & Godley, M. (1982). Alcoholism treatment by disulfiram and community reinforcement therapy. *Journal of Behavior Therapy and Experimental Psychiatry*, *13*, 105-112.
- 312 McCrady, B.S., Noel, N.E., Abrams, D.B., et al. (1986).

- 313 O'Farrell, T.J., Cutter, H.S.G., & Floyd, F.J. (1985).
- 314 Barber, J.G., & Crisp, B.R. (1995). The "pressures to change" approach to working with the partners of heavy drinkers. *Addiction, 90*(2), 269–276.
- 315 O'Farrell, T.J., Choquette, K.A., Cutter, H.S., et al. (1993).
- 316 Higgins, S.T., Budney, A.J., Bickel, W.K., et al. (1994).
- 317 Hser, Y.I., Polinsky, M.L., Maglione, M., & Anglin, M.D. (1999). Matching clients' needs with drug treatment services. *Journal of Substance Abuse Treatment, 16*, 299–305.
- 318 McLellan, A.T., Grissom, G.R., Brill, P., Durell, J., Metzger, D.S., & O'Brien, C.P. (1993). Private substance abuse treatments: Are some programs more effective than others? *Journal of Substance Abuse Treatment, 10*(3), 243–254.
- 319 McLellan, A.T., Alterman, A.I., Metzger, D.S., Grissom, G.R., Woody, G.E., Luborsky, L., et al. (1994). Similarity of outcome predictors across opiate, cocaine, and alcohol treatments: Role of treatment services. *Journal of Consulting and Clinical Psychology, 62*(6), 1141–1158.
- 320 Pringle, J.L., Edmondston, L.A., Holland, C.L., Kirisci, L., Emptage, N., Balavage, V.K., et al. (2002). The role of wrap around services in retention and outcome in substance abuse treatment: Findings from the Wrap Around Services Impact Study. *Addictive Disorders and their Treatment, 1*(4), 109–118.
- 321 Devine, J.A., & Wright, J.D. (1997). Losing the housing game. The leveling effects of substance abuse. *American Journal of Orthopsychiatry, 67*, 618–631.
- 322 Brecht, M.L., Anglin, M.D., & Wang, J.C. (1993). Treatment effectiveness for legally coerced versus voluntary methadone maintenance clients. *American Journal of Drug and Alcohol Abuse, 19*, 89–106.
- 323 DeLeon, G. (1988). Legal pressure in therapeutic communities. *Journal of Drug Issues, 18*, 625–640.
- 324 Hser, Y.I., Polinsky, M.L., Maglione, M., et al. (1999).
- 325 Pringle, J.L., Edmondston, L.A., Holland, C.L., et al. (2002).
- 326 Friedmann, P.D., Hendrickson, J.C., Gerstein, D.R., & Zhang, Z. (2004). Designated case managers as facilitators of medical and psychosocial service delivery in addiction treatment programs. *Journal of Behavior Health Services Research, 31*, 86–97.
- 327 Hser, Y.I., Polinsky, M.L., Maglione, M., et al. (1999).
- 328 McLellan, A.T., Grissom, G.R., Brill, P., et al. (1993).
- 329 Pringle, J.L., Edmondston, L.A., Holland, C.L., et al. (2002).
- 330 Friedmann, P.D., Hendrickson, J.C., Gerstein, D.R., et al. (2004).
- 331 Kessler, R.C., Berglund, P., Demler, O., Jin, R., Merikangas, K.R., & Walters, E.E. (2005). Lifetime prevalence and age-of-onset distributions of DSM-IV disorders in the National Comorbidity Survey Replication. *Archives of General Psychiatry, 62*, 593–602.
- 332 Ross, H.E., Glaser, F.B., & Germanson, T. (1988). The prevalence of psychiatric disorders in patients with alcohol and other drug problems. *Archives of General Psychiatry, 45*, 1023–1031.
- 333 Anglin, M.D., Hser, Y.I., & Booth, M.W. (1987). Sex differences in addict careers. 4. Treatment. *American Journal of Drug and Alcohol Abuse, 13*, 253–280.
- 334 Grella, C.E., & Joshi, V. (1999). Gender differences in drug treatment careers among clients in the National Drug Abuse Treatment Outcome Study. *American Journal of Drug and Alcohol Abuse, 25*(3), 385–406.
- 335 Wechsberg, W.M., Craddock, S.G., & Hubbard, R.L. (1998). How are women who enter substance abuse treatment different than men? A gender comparison from the Drug Abuse Treatment Outcome Study (DATOS). *Drugs and Society, 13*(1/2), 97–115.
- 336 Tims, F., Leukefeld, C., & Platt, J. (2001). *Relapse and recovery in addictions*. New Haven, CT: Yale University Press.
- 337 Halfon, N., & Hochstein, M. (2002). Life course health development: An integrated framework for developing health, policy, and research. *Milbank Quarterly, 80*, 433–79, iii.
- 338 Antonucci, T.C. (1990). Social supports and social relationships. In R.H. Binstock (Ed.), *Handbook of aging and the social sciences* (3rd ed.). New York: Van Nostrand Reinhold.
- 339 Rice, C., Longabaugh, R., Beattie, M., & Noel, N. (1993). Age group differences in response to treatment for problematic alcohol use. *Addiction, 88*(10), 1369–1375.
- 340 Bartels, S.J., Blow, F.C., Brockmann, L.M., & Van Citters, A.D. (2005). *Substance abuse and mental health among older Americans: The state of the knowledge and future directions*. Rockville, MD: Westat.

- 341 Blow, F.C., Walton, M.A., Chermack, S.T., Mudd, S.A., & Brower, K.J. (2000). Older adult treatment outcome following elder-specific inpatient alcoholism treatment. *Journal of Substance Abuse Treatment, 19*(1), 67–75.
- 342 Schonfeld, L., & Dupree, L.W. (1995). Treatment approaches for older problem drinkers. *International Journal of the Addictions, 30*(13–14), 1819–1842.
- 343 Benschhoff, J.J., Harrawood, L.K., & Koch, D.S. (2003). Substance abuse and the elderly: Unique issues and concerns. *Journal of Rehabilitation, 69*, 43–48.
- 344 Satre, D.D., Mertens, J.R., Areal, P.A., & Weisner, C. (2004). Five-year alcohol and drug treatment outcomes of older adults versus middle-aged and younger adults in a managed care program. *Addiction, 99*, 1286–1297.
- 345 Tims, F., Leukefeld, C., & Platt, J. (2001).
- 346 Blow, F.C., Walton, M.A., Chermack, S.T., et al. (2000).
- 347 Schonfeld, L., & Dupree, L.W. (1995).
- 348 Benschhoff, J.J., Harrawood, L.K., & Koch, D.S. (2003).
- 349 Berk, L.E. (2007). *Development through the lifespan* (4th ed.). Boston: Allyn & Bacon.
- 350 Featherman, D.L., & Lerner, R.M. (1985). Ontogenesis and sociogenesis: Problematics for theory and research about development and socialization across the lifespan. *American Sociological Review, 50*(5), 659–676.
- 351 Rice, C., Longabaugh, R., Beattie, M., et al. (1993).
- 352 Schonfeld, L., & Dupree, L.W. (1995).
- 353 Bartels, S.J., Blow, F.C., Brockmann, L.M., et al. (2005).
- 354 Blow, F.C., Walton, M.A., Chermack, S.T., et al. (2000).
- 355 Fortney, J.C., Booth, B.M., Blow, F.C., & Bunn, J.Y. (1995). The effects of travel barriers and age on the utilization of alcoholism treatment aftercare. *American Journal of Drug and Alcohol Abuse, 21*, 391–406.
- 356 Rice, C., Longabaugh, R., Beattie, M., et al. (1993).
- 357 Bartels, S.J., Blow, F.C., Brockmann, L.M., et al. (2005).
- 358 Blow, F.C., Walton, M.A., Chermack, S.T., et al. (2000).
- 359 Schonfeld, L., & Dupree, L.W. (1995).
- 360 Fillmore, K.M., Grant, M., Hartka, E., et al. (1988).
- 361 Sobell, L.C., Ellingstad, T.P., & Sobell, M.B. (2000).
- 362 Humphreys, K., Moos, R.H., & Finney, J.W. (1995). Two pathways out of drinking problems without professional treatment. *Addictive Behaviors, 20*, 427–441.
- 363 Granfield, R., & Cloud, W. (2001).
- 364 Drake, R.E., Wallach, M.A., & McGovern, M.P. (2005).
- 365 Hser, Y.I., Grella, C.E., Hsieh, S.C., Anglin, M.D., & Brown, B.S. (1999). Prior treatment experience related to process and outcomes in DATOS. *Drug and Alcohol Dependence, 57*, 137–150.
- 366 Simpson, D.D., Joe, G.W., Greener, J.M., & Rowan-Szal, G.A. (2000). Modeling year 1 outcomes with treatment process and post-treatment social influences. *Substance Use and Misuse, 35*, 1911–1930.
- 367 Broome, K.M., Simpson, D.D., & Joe, G.W. (2002). The role of social support following short-term inpatient treatment. *American Journal of Addictions, 11*, 57–65.
- 368 Bradizza, C.M., & Stasiewicz, P.R. (2003). Qualitative analysis of high-risk drug and alcohol use situations among severely mentally ill substance abusers. *Addictive Behaviors, 28*, 157–169.
- 369 Bradizza, C.M., Stasiewicz, P.R., & Carey, K.B. (1998). High-risk alcohol and drug use situations among seriously mentally ill inpatients: A preliminary investigation. *Addictive Behaviors, 23*, 555–560.
- 370 Alverson, H., Alverson, M., & Drake, R.E. (2000). An ethnographic study of the longitudinal course of substance abuse among people with severe mental illness. *Community Mental Health Journal, 36*, 557–569.
- 371 Sells, D.J., Rowe, M., Fisk, D., & Davidson, L. (2003). Violent victimization of persons with co-occurring psychiatric and substance use disorders. *Psychiatric Services, 54*, 1253–1257.
- 372 Sells, D.J., Rowe, M., Fisk, D., et al. (2003).
- 373 Kisthardt, W. (1997). The strengths model of case management: Principles and helping functions. In D. Saleebey (Ed.), *The Strengths Perspective in Social Work Practice* (2nd ed., pp. 97–114). New York: Longman.

- 374 Rapp, C.A. (1998). *The strengths model: Case management with people suffering from severe and persistent mental illness*. New York: Oxford University Press.
- 375 Carten, A.J. (1996).
- 376 Rapp, C.A. (1998).
- 377 Blankertz, L., McKay, C., & Robinson, S. (1998). Work as a rehabilitative tool for individuals with dual diagnosis. *Journal of Vocational Rehabilitation, 11*, 113–123.
- 378 Comerford, A.W. (1999). Work dysfunction and addiction: Common roots. *Journal of Substance Abuse Treatment, 16*, 247–253.
- 379 Dickinson, K., & Maynard, R. (1981). *The impact of supported work on ex-addicts*. New York: MDRC.
- 380 Platt, J.J. (1995). Vocational rehabilitation of drug abusers. *Psychological Bulletin, 117*, 416–433.
- 381 Room, R. (1998). Drinking patterns and alcohol-related social problems: Frameworks for analysis in developing societies. *Drug and Alcohol Review, 17*, 389–398.
- 382 Pringle, J.L., Edmondston, L.A., Holland, C.L., et al. (2002).
- 383 McDaid, D., & Thornicroft, G. (2005). *Mental health II: Balancing institutional and community-based care*. World Health Organization Regional Office for Europe.
- 384 Trieman, N., Leff, J., & Glover, G. (1999). Outcome of long stay psychiatric patients resettled in the community: Prospective cohort study. *British Medical Journal, 319*, 13–16.
- 385 Ibid.
- 386 Harrison, L.D. (2001). The revolving prison door for drug involved offenders: Challenges and opportunities. *Crime & Delinquency, 47*(3), 462–485.
- 387 Knight, K., Simpson, D.D., & Hiller, M. (1999). Three-year reincarceration outcomes for in-prison therapeutic community treatment in Texas. *Prison Journal, 79*, 337–351.
- 388 Haggerty, J.L., Reid, R.J., Freeman, G.K., Starfield, B.H., Adair, C.E., & McKendry, R. (2003). Continuity of care: A multidisciplinary review. *British Medical Journal, 327*, 1219–1221.
- 389 Ibid.
- 390 Moos, R.H. (2003). Addictive disorders in context: Principles and puzzles of effective treatment and recovery. *Psychology of Addictive Behaviors, 17*, 3–12.
- 391 Miller, W.R., & Hester, R.K. (1986). Inpatient alcoholism treatment: Who benefits? *American Psychologist, 41*, 794–805.
- 392 Fortney, J.C., Booth, B.M., Blow, F.C., et al. (1995).
- 393 McKay, J.R. (2001). Effectiveness of continuing care interventions for substance abusers: Implications for study of long-term treatment effects. *Evaluation Review, 25*(2), 211–232.
- 394 Gruber, K.J., Fleetwood, T.W., & Herring, M.W. (2001).
- 395 Gruber, K.J., & Fleetwood, T.W. (2004).
- 396 Godley, M.D., Godley, S.H., Dennis, M.L., Funk, R., & Passetti, L.L. (2002). Preliminary outcomes from the assertive continuing care experiment for adolescents discharged from residential treatment. *Journal of Substance Abuse Treatment, 23*, 21–32.
- 397 Dennis, M., Scott, C.K., & Funk, R. (2003). An experimental evaluation of recovery management checkups (RMC) for people with chronic substance use disorders. *Evaluation and Program Planning, 26*, 339–352.
- 398 Donovan, D.M. (1998).
- 399 Godley, M.D., Godley, S.H., Dennis, M., Funk, R., & Passetti, L.L. (2007). The effect of assertive continuing care on continuing care linkage, adherence and abstinence following residential treatment for adolescents with substance use disorders. *Addiction, 102*, 81–93.
- 400 Godley, M.D., Godley, S.H., Dennis, M., et al. (2007).
- 401 Druss, B.G., & von Esenwein, S.A. (2006). Improving general medical care for persons with mental and addictive disorders: Systematic review. *General Hospital Psychiatry, 28*, 145–153.
- 402 Rothman, J., Rudnick, D., Slifer, M., Agins, B., Heiner, K., & Birkhead, G. (2007). Co-located substance use treatment and HIV prevention and primary care services, New York State, 1990-2002: A model for effective service delivery to a high-risk population. *Journal of Urban Health, 84*, 226–242.
- 403 Hitchcock, H.C., Stainback, R.D., & Roque, G.M. (1995). Effects of halfway house placement on retention of patients in substance abuse aftercare. *American Journal of Drug and Alcohol Abuse, 21*, 379–390.

- 404 Rosenheck, R., & Gallup, P. (1991). Involvement in an outreach and residential treatment program for homeless mentally ill veterans. *Journal of Nervous and Mental Disease, 179*, 750–754.
- 405 Moos, R.H., Finney, J.W., Ouimette, P.C., & Suchinsky, R.T. (1999). A comparative evaluation of substance abuse treatment: I. Treatment orientation, amount of care, and 1-year outcomes. *Alcoholism, Clinical and Experimental Research, 23*, 529–536.
- 406 Farabee, D., Prendergast, M., Cartier, J., Wexler, H., Knight, K., & Anglin, M.D. (1999). Barriers to implementing effective correctional drug treatment programs. *Prison Journal, 7*(2), 150–162.
- 407 Wolff, N., Plemmons, D., Veysey, B., & Brandli, A. (2002). Release planning for inmates with mental illness compared with those who have other chronic illnesses. *Psychiatric Services, 53*, 1469–1471.
- 408 Sung, H.E., & Richter, L. (2006). Contextual barriers to successful reentry of recovering drug offenders. *Journal of Substance Abuse Treatment, 31*, 365–374.
- 409 Butzin, C.A., Martin, S.S., & Inciardi, J.A. (2002). Evaluating component effects of a prison-based treatment continuum. *Journal of Substance Abuse Treatment, 22*, 63–69.
- 410 Wexler, H.K., Melnick, G., Lowe, L., & Peters, J. (1999). Three-year reincarceration outcomes for Amity in-prison therapeutic community and aftercare in California. *Prison Journal, 79*, 321–336.
- 411 Broome, K.M., Simpson, D.D., & Joe, G.W. (1999). Patient and program attributes related to treatment process indicators in DATOS. *Drug and Alcohol Dependence, 57*, 127–135.
- 412 Joe, G.W., Simpson, D.D., Dansereau, D.F., & Rowan-Szal, G.A. (2001). Relationships between counseling rapport and drug abuse treatment outcomes. *Psychiatric Services, 52*, 1223–1229.
- 413 Alverson, H., Alverson, M., & Drake, R.E. (2001). Social patterns of substance use among people with dual diagnoses. *Mental Health Services Research, 3*, 3–14.
- 414 Moos, R.H., Finney, J.W., Ouimette, P.C., et al. (1999).
- 415 Farabee, D., Prendergast, M., Cartier, J., et al. (1999).
- 416 Wolff, N., Plemmons, D., Veysey, B., et al. (2002).
- 417 Sung, H.E., & Richter, L. (2006).
- 418 Butzin, C.A., Martin, S.S., & Inciardi, J.A. (2002).
- 419 Wexler, H.K., Melnick, G., Lowe, L., et al. (1999).
- 420 Broome, K.M., Simpson, D.D., & Joe, G.W. (1999).
- 421 Ibid,
- 422 Joe, G.W., Simpson, D.D., Dansereau, D.F., et al. (2001).
- 423 Alverson, H., Alverson, M., & Drake, R.E. (2001).
- 424 Green, J.M. (1996). Engagement and empathy: A pilot study of the therapeutic alliance in outpatient child psychiatry. *Child Psychology and Psychiatry Review, 1*, 130–138.
- 425 Meier, P.S., Barrowclough, C., & Donmall, M.C. (2005). The role of the therapeutic alliance in the treatment of substance misuse: A critical review of the literature. *Addiction, 100*, 304–316.
- 426 Felton, B.J., Barr, A., Clark, G., & Tsemberis, S.J. (2006). ACT team members' responses to training in recovery-oriented practices. *Psychiatric Rehabilitation Journal, 30*, 112–119.
- 427 Lorig, K.R., & Holman, H. (2003).
- 428 Ibid, p. 2.
- 429 Ibid, p. 1.
- 430 Moos, R.H. (2003).
- 431 Simpson, D.D. (2004). A conceptual framework for drug treatment process and outcomes. *Journal of Substance Abuse Treatment, 27*, 99–121.
- 432 Connors, G.J., Carroll, K.M., DiClemente, C.C., Longabaugh, R., & Donovan, D.M. (1997). The therapeutic alliance and its relationship to alcoholism treatment participation and outcome. *Journal of Consulting and Clinical Psychology, 65*, 588–598.
- 433 Dearing, R.L., Barrick, C., Dermen, K.H., & Walitzer, K.S. (2005). Indicators of client engagement: Influences on alcohol treatment satisfaction and outcomes. *Psychology of Addictive Behaviors, 19*, 71–78.
- 434 Miller, W.R., & Wilbourne, P.L. (2002). Mesa Grande: A methodological analysis of clinical trials of treatments for alcohol use disorders. *Addiction, 97*, 265–277.
- 435 Dearing, R.L., Barrick, C., Dermen, K.H., et al. (2005).

- 436 Belding, M.A., Iguchi, M.Y., Morral, A.R., & McLellan, A.T. (1997). Assessing the helping alliance and its impact in the treatment of opiate dependence. *Drug and Alcohol Dependence*, 48, 51–59.
- 437 Petry, N.M., & Bickel, W.K. (1999). Therapeutic alliance and psychiatric severity as predictors of completion of treatment for opioid dependence. *Psychiatric Services*, 50, 219–227.
- 438 Diamond, G.M., Diamond, G.S., & Liddle, H.A. (2000). The therapist-parent alliance in family-based therapy for adolescents. *Journal of Clinical Psychology*, 56, 1037–1050.
- 439 Kasarabada, N.D., Hser, Y.I., Boles, S.M., & Huang, Y.C. (2002). Do patients' perceptions of their counselors influence outcomes of drug treatment? *Journal of Substance Abuse Treatment*, 23(4), 327–334.
- 440 Ilgen, M., Tiet, Q., Finney, J., & Moos, R.H. (2006). Self-efficacy, therapeutic alliance, and alcohol-use disorder treatment outcomes. *Journal of Studies on Alcohol*, 67(3), 465–472.
- 441 Ilgen, M.A., McKellar, J., Moos, R., & Finney, J.W. (2006). Therapeutic alliance and the relationship between motivation and treatment outcomes in patients with alcohol use disorder. *Journal of Substance Abuse Treatment*, 31(2), 157–162. Epub 2006 Jul 13.
- 442 Carten, A.J. (1996).
- 443 Siegal, H.A., Fisher, J.H., Rapp, R.C., Kelliher, C.W., Wagner, J.H., O'Brien, W.F., et al. (1996). Enhancing substance abuse treatment with case management: Its impact on employment. *Journal of Substance Abuse Treatment*, 13(2), 93–98.
- 444 Hall, J.A., Carswell, C., Walsh, E., Huber, D.L., & Jampoler, J.S. (2002). Iowa case management: Innovative social casework. *Social Work*, 47, 132–141.
- 445 Rapp, R.C., Kelliher, C.W., Fisher, J.H., & Hall, F.J. (1994). Strengths-based case management. A role in addressing denial in substance abuse treatment. *Journal of Case Management*, 3, 139–144.
- 446 Saleebey, D. (1996). The strengths perspective in social work practice: Extensions and cautions. *Social Work*, 41, 296–305.
- 447 Weick, A., Rapp, C., Sullivan, W.P., & Kisthardt, W. (1989). A strengths perspective for social work practice. *Social Work*, 34 (4): 350–354.
- 448 Siegal, H.A., Fisher, J.H., Rapp, R.C., et al. (1996).
- 449 Siegal, H.A., Li, L., & Rapp, R.C. (2002). Case management as a therapeutic enhancement: Impact on post-treatment criminality. *Journal of Addictive Disorders*, 21(4), 37–46.
- 450 Rapp, R.C., Siegal, H.A., Li, L., & Saha, P. (1998). Predicting post primary treatment services and drug use outcome: A multivariate analysis. *American Journal of Drug and Alcohol Abuse*, 24, 603–615.
- 451 Arredondo, P., Toperek, R., Brown, S.P., Jones, J., Locke, D.C., Sanchez, J., et al. (1996). Operationalization of the multicultural counseling competencies. *Journal of Multicultural Counseling and Development*, 24, 42–78.
- 452 Office of Minority Health (2001). *National standards for culturally and linguistically appropriate services in health care: Final report*. Rockville, MD: U.S. Department of Health and Human Services. Retrieved April 2, 2007, from <http://www.omhrc.gov/templates/browse.aspx?lvl=2&lvlID=15>.
- 453 Brach, C., & Fraser, I. (2000). Can cultural competency reduce racial and ethnic health disparities? A review and conceptual model. *Medical Care Research Review*, 57 (S1): 181–217.
- 454 Lavizzo-Mourey, R., & Mackenzie, E.R. (1996). Cultural competence: Essential measurements of quality for managed care organizations. *Annals of Internal Medicine*, 124 (10): 919–920.
- 455 Lawson, W.B. (1996). The art and science of the psychopharmacotherapy of African Americans. *Mount Sinai Journal of Medicine*, 63(5–6): 301–305.
- 456 Moffic, H.S., & Kinzie, J.D. (1996). The history and future of cross-cultural psychiatric services. *Community Mental Health Journal*, 32(6): 581–592.
- 457 Cheung, F.K., & Snowden, L.R. (1990). Community mental health and ethnic minority populations. *Community Mental Health Journal*, 26, 277–291.
- 458 Simpson, D.D. (2004).
- 459 Office of Minority Health (2001).
- 460 Brach, C., & Fraser, I. (2000).
- 461 Ford, E.S., & Cooper, R.S. (1995). Implications of race/ethnicity for health and health care use. *Health Services Research*, 30(1), 237–252.

- 462 Gornick, M.E., Eggers, P.W., Reilly, T.W., Mentnech, R.M., Fitterman, L.K., Kucken, L.E., et al. (1996). Effects of race and income on mortality and use of services among Medicare beneficiaries. *New England Journal of Medicine*, 335(11), 791–799.
- 463 Mayberry, R.M., Mili, F., Vaid, I.G.M., Samadi, A., Ofili, E., McNeal, M.S., et al. (1999). *Racial and ethnic differences in access to medical care: A synthesis of the literature*. Menlo Park, CA: Henry J. Kaiser Family Foundation.
- 464 Kington, R.S., & Smith, J.P. (1997). Socioeconomic status and racial and ethnic differences in functional status associated with chronic diseases. *American Journal of Public Health*, 87(5), 805–810.
- 465 Williams, D.R. (1999). Race, socioeconomic status, and health: The added effects of racism and discrimination. *Annals of the New York Academy of Sciences*, 896, 173–188.
- 466 Nickens, H.W. (1995). The role of race/ethnicity and social class in minority health status. *Health Services Research*, 30 (1, Part II), 151–162.
- 467 Hyun, M.S., Kools, S., & Kim, S.A. (2003).
- 468 Uziel-Miller, N.D., Lyons, J.S., Kissiel, C., & Love, S. (1998). Treatment needs and initial outcomes of a residential recovery program for African-American women and their children. *American Journal of Addictions*, 7, 43–50.
- 469 Campbell, C.I., & Alexander, J.A. (2002). Culturally competent treatment practices and ancillary service use in outpatient substance abuse treatment. *Journal of Substance Abuse Treatment*, 22(3), 109–119.
- 470 Longshore, D., Grills, C., & Annon, K. (1999). Effects of a culturally congruent intervention on cognitive factors related to drug-use recovery. *Substance Use and Misuse*, 34(9), 1223–1241.
- 471 Campbell, C.I., & Alexander, J.A. (2002).
- 472 Ehrmin, J.T. (2005). Dimensions of culture care for substance-dependent African American women. *Journal of Transcultural Nursing*, 16, 117–125.
- 473 Miller, W.R. (1998).
- 474 Miller, W.R., Meyers, R.J., & Hiller-Sturmhofel, S. (1999). The community-reinforcement approach. *Alcohol Research and Health*, 23, 116–121.
- 475 Ehrmin, J.T. (2005).
- 476 Arnold, R., Avants, S.K., Margolin, A., & Marcotte, D. (2002). Patient attitudes concerning the inclusion of spirituality into addiction treatment. *Journal of Substance Abuse Treatment*, 23, 319–326.
- 477 Avants, S.K., Warburton, L.A., & Margolin, A. (2001). Spiritual and religious support in recovery from addiction among HIV-positive injection drug users. *Journal of Psychoactive Drugs*, 33, 39–45.
- 478 Brizer, D.A. (1993). Religiosity and drug abuse among psychiatric inpatients. *American Journal of Drug and Alcohol Abuse*, 19, 337–345.
- 479 Gorsuch, R.L. (1994). Toward motivational theories of intrinsic religious commitment. *Journal of Science and Studies in Religion*, 33(4), 315–325.
- 480 Kendler, K.S., Gardner, C.O., & Prescott, C.A. (1997). Religion, psychopathology, and substance use and abuse: A multimeasure, genetic-epidemiologic study. *American Journal of Psychiatry*, 154, 322–329.
- 481 Mathew, R.J., Georgi, J., Wilson, W.H., & Mathew, V.G. (1996). A retrospective study of the concept of spirituality as understood by recovering individuals. *Journal of Substance Abuse Treatment*, 13, 67–73.
- 482 Miller, W.R. (1998).
- 483 Avants, S.K., Warburton, L.A., & Margolin, A. (2001).
- 484 Muffler, J., Langrod, J.G., & Larson, D. (1992). “There is a balm in Gilead.” Religion and substance abuse treatment. In J.H. Lowinson, P. Ruiz, & R.B. Millman (Eds.), *Substance abuse: A comprehensive textbook* (2nd ed., pp. 584–595). Baltimore: Williams & Wilkins.
- 485 Laudet, A. B., Morgen, K., & White, W. L. (2006).
- 486 Ibid.
- 487 Pardini, D.A., Plante, T.G., Sherman, A., & Stump, J.E. (2000). Religious faith and spirituality in substance abuse recovery: Determining the mental health benefits. *Journal of Substance Abuse Treatment*, 19, 347–354.
- 488 Kondo, C., Iimuro, T., Iwai, K., Kurata, K., Kouda, M., Tachikawa, H., et al. (2000). [A study of recovery factor about drug addiction recovery center (DARC)]. *Nihon Arukoru Yakubutsu Igakkai Zasshi*, 35, 258–270.
- 489 Flynn, P.M., Joe, G.W., Broome, K.M., et al. (2003).

- 490 Laudet, A. B., Morgen, K., & White, W. L. (2006). p. 15.
- 491 Ibid.
- 492 Miller, W.R. (1998). Why do people change addictive behavior? The 1996 H. David Archibald Lecture. *Addiction*, *93*, 163–172.
- 493 Arnold, R., Avants, S.K., Margolin, A., et al. (2002).
- 494 Institute of Medicine. (2006).
- 495 Humphreys, K., Wing, S., McCarty, D., et al. (2004).
- 496 Galanter, M., Dermatis, H., Egelko, S., & De Leon, G. (1998). Homelessness and mental illness in a professional- and peer-led cocaine treatment clinic. *Psychiatric Services*, *49*(4), 533–535.
- 497 Bandura, A. (1997). The anatomy of stages of change. *American Journal of Health Promotion*, *12*, 8–10.
- 498 Lorig, K.R., Ritter, P., Stewart, A.L., et al. (2001).
- 499 Institute of Medicine. (2006).
- 500 Bodenheimer, T., Lorig, K., Holman, H., & Grumbach, K. (2002). Patient self-management of chronic disease in primary care. *JAMA*, *288*, 2469–2475.
- 501 Moos, R.H., Moos, B.S., Timko, C. (2006). Gender, Treatment, and Self-Help in Remission from Alcohol Use and Disorders. *Clinical Medical Research*, *4*, 163-174.
- 502 Morgenstern, J., Labouvie, E., McCrady, B.S. et al. (1997).
- 503 Humphreys, K. (2004). A few apologies, but no regrets. *Addiction*, *99*, 155–156.
- 504 Humphreys, K., Mankowski, E., Moos, R., & Finney, J. (1999). Do enhanced friendship networks and active coping mediate the effect of self-help groups on substance abuse? *Annals of Behavioral Medicine*, *21*(1), 54–60.
- 505 McKellar, J., Stewart, E., & Humphreys, K. (2003). Alcoholics anonymous involvement and positive alcohol-related outcomes: Cause, consequence, or just a correlate? A prospective 2-year study of 2,319 alcohol-dependent men. *Journal of Consulting and Clinical Psychology*, *71*, 302–308.
- 506 Connors, G.J., Tonigan, J.S., & Miller, W.R. (2001). A longitudinal model of intake symptomatology, AA participation and outcome: Retrospective study of the project MATCH outpatient and aftercare samples. *Journal of Studies on Alcohol*, *62*, 817–825.
- 507 Owen, P.L., Slaymaker, V., Tonigan, J.S., McCrady, B.S., Epstein, E.E., Kaskutas, L.A., et al. (2003). Participation in Alcoholics Anonymous: Intended and unintended change mechanisms. *Alcoholism, Clinical and Experimental Research*, *27*, 524–532.
- 508 Humphreys, K., & Moos, R. (2001). Can encouraging substance abuse patients to participate in self-help groups reduce demand for health care? A quasi-experimental study. *Alcoholism, Clinical and Experimental Research*, *25*, 711–716.
- 509 Hawkins, J.D., & Catalano, R.F. (1985). Aftercare in drug abuse treatment. *International Journal of the Addictions*, *20*, 917–945.
- 510 Johnson, N.P., & Chappel, J.N. (1994). Using AA and other 12-step programs more effectively. *Journal of Substance Abuse Treatment*, *11*(2), 137–142.
- 511 Khantzian, E.J., & Mack, J.E. (1994). How AA works and why it's important for clinicians to understand. *Journal of Substance Abuse Treatment*, *11*(2), 77–92.
- 512 Connors, G.J., Tonigan, J.S., & Miller, W.R. (2001).
- 513 Humphreys, K., Mankowski, E., Moos, R., et al. (1999).
- 514 Owen, P.L., Slaymaker, V., Tonigan, J.S., et al. (2003).
- 515 Humphreys, K., & Moos, R. (2001).
- 516 Johnson, N.P., & Chappel, J.N. (1994).
- 517 Khantzian, E.J., & Mack, J.E. (1994).
- 518 Hawkins, J.D., & Catalano, R.F. (1985).
- 519 Troyer, T.N., Acampora, A.P., O'Connor, L.E., & Berry, J.W. (1995). The changing relationship between therapeutic communities and 12-Step programs: A survey. *Journal of Psychoactive Drugs*, *27*, 177–180.
- 520 Fiorentine, R., & Hillhouse, M.P. (2000).
- 521 Caldwell, P.E., & Cutter, H.S. (1998). Alcoholics Anonymous affiliation during early recovery. *Journal of Substance Abuse Treatment*, *15*, 221–228.

- 522 Etheridge, R.M., Craddock, S.G., Hubbard, R.L., & Rounds-Bryant, J.L. (1999). The relationship of counseling and self-help participation to patient outcomes in DATOS. *Drug and Alcohol Dependence*, 57, 99–112.
- 523 Vanicelli, M. (1978). Impact of aftercare in the treatment of alcoholics: A cross-lagged panel analysis. *Journal of Studies on Alcohol*, 39, 1875–1886.
- 524 Kissina, W., McLeoda, C., & McKay, J. (2003). The longitudinal relationship between self-help group attendance and course of recovery. *Evaluation and Program Planning*, 26, 311–323.
- 525 Morgenstern, J., Labouvie, E., McCrady, B.S., et al. (1997).
- 526 Humphreys, K., Mankowski, E., Moos, R., et al. (1999).
- 527 Galanter, M., Egelko, S., & Edwards, H. (1993). Rational recovery: Alternative to AA for addiction? *American Journal of Drug and Alcohol Abuse*, 19, 499–510.
- 528 Brooks, A.J., & Penn, P.E. (2003). Comparing treatments for dual diagnosis: Twelve-step and self-management and recovery training. *American Journal of Drug and Alcohol Abuse*, 29, 359–383.
- 529 Scutchfield, F.D., Ireson, C., & Hall, L. (2004). The voice of the public in public health policy and planning: The role of public judgment. *Journal of Public Health Policy*, 25, 197–205.
- 530 Merzel, C., & D’Afflitti, J. (2003). Reconsidering community-based health promotion: Promise, performance, and potential. *American Journal of Public Health*, 93(4), 557–574.
- 531 Ibid.
- 532 Jeffs, L., Law, M., Baker, G.R., & Norton, P.G. (2005). *Patient safety research in Australia, United Kingdom, United States and Canada: A summary of research priority areas, agenda-setting processes, and directions for future research in the context of their patient safety initiatives*. Canadian Patient Safety Institute.
- 533 Ibid.
- 534 Dwyer, J. (1989). The politics of participation. *Community Health Studies*, 13, 59–65.
- 535 Sofaer, S. (1999). Challenges for the public in negotiating the health system in the 21st century. *Journal of Urban Health*, 76, 211–228.
- 536 Walsh, D. (1996). A journey toward recovery: From the inside out. *Psychiatric Rehabilitation Journal*, 20(2), 85–90.
- 537 Sofaer, S. (1999).
- 538 Ibid.
- 539 Moggi, F., Hirsbrunner, H.P., Brodbeck, J., & Bachmann, K.M. (1999). One-year outcome of an integrative inpatient treatment for dual diagnosis patients. *Addictive Behaviors*, 24(4), 589–592.
- 540 Weisner, C., Mertens, J., Parthasarathy, S., Moore, C., & Lu, Y. (2001). Integrating primary medical care with addiction treatment: A randomized controlled trial. *JAMA*, 286, 1715–1723.
- 541 Parthasarathy, S., Mertens, J., Moore, C., & Weisner, C. (2003). Utilization and cost impact of integrating substance abuse treatment and primary care. *Medical Care*, 41, 357–367.
- 542 McAlpine, C., Marshall, C.C., & Doran, N.H. (2001). Combining child welfare and substance abuse services: A blended model of intervention. *Child Welfare*, 80, 129–149.
- 543 Friedmann, P.D., Hendrickson, J.C., Gerstein, D.R., et al. (2004).
- 544 Weisner, C., Mertens, J., Parthasarathy, S., et al. (2001).
- 545 Friedmann, P.D., Lemon, S.C., Stein, M.D., Etheridge, R.M., D’Aunno, T. A. (2001). Linkage to medical services in the Drug Abuse Treatment Outcome Study. *Medical Care*, 39, 284–295.
- 546 Parthasarathy, S., Mertens, J., Moore, C., et al. (2003).
- 547 McAlpine, C., Marshall, C.C., & Doran, N.H. (2001)
- 548 McLellan, A.T., Hagan, T.A., Levine, M., Gould, F., Meyers, K., Bencivengo, M., et al. (1998). Supplemental social services improve outcomes in public addiction treatment. *Addiction*, 93, 1489–1499.
- 549 Marsh, J.C., D’Aunno, T.A., & Smith, B.D. (2000). Increasing access and providing social services to improve drug abuse treatment for women with children. *Addiction*, 95, 1237–1247.
- 550 Minkoff, K. (1991). Program components of a comprehensive integrated care system for serious mentally ill patients with substance disorders. *New Directions in Mental Health Services*, (50), 13–27.
- 551 Osher, F.C., & Kofoed, L.L. (1989). Treatment of patients with psychiatric and psychoactive substance abuse disorders. *Hospital and Community Psychiatry*, 40, 1025–1030.

- 552 Swindle, R.W., Phibbs, C.S., Paradise, M.J., Recine, B.P., & Moos, R.H. (1995). Inpatient treatment for substance abuse patients with psychiatric disorders: A national study of determinants of readmission. *Journal of Substance Abuse*, 7(1), 79–97.
- 553 Moggi, F., Hirsbrunner, H.P., Brodbeck, J., et al. (1999)
- 554 Weisner, C., Mertens, J., Parthasarathy, S., et al. (2001).
- 555 Golden, O. (1992). *Poor children and welfare reform*. Westport, CT: Auburn House.
- 556 Swick, K.J. (1992). *A descriptive assessment of project FOCUS' home visit program*. (ERIC Document Reproduction Service No. ED 346962)
- 557 Boyd, B. (1992). Impacts of interagency collaboration on participating organizations. Paper presented at the annual meeting of the American Educational Research Association, San Francisco, CA: (ERIC Document Reproduction Service No. ED 346566)
- 558 Kagan, S.L. (1991). *United we stand: Collaboration for childcare and early education services*. New York: Teachers College Press.
- 559 Raack, L., Kunesh, L.G., & Shulman, I. (1992). *Interagency collaboration in the heartland: Challenges and opportunities. Proceedings of the NCREL Early Childhood Connection Regional Forum (1st, Elmhurst, Illinois, October 14-15, 1991)*. ERIC Document Reproduction Service No. ED 344670.
- 560 Washburn, N.J., Simmons, S., Sommer, V., Adkins, B., Gerken, P., Rogers, M., et al. (2006). Outcome of interventions to identify family history and risk management for women with breast cancer in the ambulatory setting. *Journal of Clinical Oncology*, 24(18S), 6121.
- 561 Weingarten, S.R., Henning, J.M., Badamgarav, E., Knight, K., Hasselblad, V., Gano, A., Jr., et al. (2002). Interventions used in disease management programmes for patients with chronic illness—which ones work? Meta-analysis of published reports. *British Medical Journal*, 325, 925.
- 562 Davis, D.A., Thomson, M.A., Oxman, A.D., Haynes, R.B. (1992). Evidence for the effectiveness of CME. A review of 50 randomized controlled trials. *JAMA*, 268, 1111–1117.
- 563 Weingarten, S.R., Henning, J.M., Badamgarav, E., et al. (2002).
- 564 Bukstein, D.A., Jones, C.A., Ledford, D.K., Smith, P., Wechsler, M.E., & Urbano, F.L. (2005). Discussing the COSTS of Asthma: Controlling Outcomes, Symptoms, and Treatment Strategies. *American Journal of Managed Care*, 11(11S), S318–S336.
- 565 Dennis, M., Scott, C.K., & Funk, R. (2003).
- 566 Dubar-Jacob, J., Burke, L.E., & Puczynski, S. (1995). Clinical assessment and management of adherence to medical regimens. In P.M. Nicassio, & T.W. Smith (Eds.), *Managing chronic illness: A biosychosocial perspective* (pp. 313–349). Washington, DC: American Psychological Association.
- 567 Nicassio, P.M., Schoenfeld-Smith, K., Radojevic, V., & Schuman, C. (1995). Pain coping mechanisms in fibromyalgia: Relationship to pain and functional outcomes. *Journal of Rheumatology*, 22, 1552–1558.
- 568 Roter, D.L., Hall, J.A., Merisca, R., Nordstrom, B., Cretin, D., & Svarstad, B. (1998). Effectiveness of interventions to improve patient compliance: A meta-analysis. *Medical Care*, 36, 1138–1161.
- 569 McLellan, A.T., Lewis, D.C., O'Brien, C.P., et al. (2000).
- 570 Dennis, M., Scott, C.K., & Funk, R. (2003).
- 571 Institute of Medicine. (2006).
- 572 McKay, J.R., Alterman, A.I., Rutherford, M.J., Cacciola, J.S., & McLellan, A.T. (1999). The relationship of alcohol use to cocaine relapse in cocaine dependent patients in an aftercare study. *Journal of Studies on Alcohol*, 60, 176–180.
- 573 McKay, J.R., Lynch, K.G., Shepard, D.S., Ratichek, S., Morrison, R., Koppenhaver, J., et al. (2004). The effectiveness of telephone-based continuing care in the clinical management of alcohol and cocaine use disorders: 12-month outcomes. *Journal of Consulting and Clinical Psychology*, 72, 967–979.
- 574 McLellan, A.T. (2002). Have we evaluated addiction treatment correctly? Implications from a chronic care perspective. *Addiction*, 97, 249–252.
- 575 Hser, Y.I., Joshi, V., Anglin, M.D., & Fletcher, B. (1999). Predicting posttreatment cocaine abstinence for first-time admissions and treatment repeaters. *American Journal of Public Health*, 89, 666–671.
- 576 Simpson, D.D., & Savage, L.J. (1980). Drug abuse treatment readmissions and outcomes: Three-year follow-up of DARP patients. *Archives of General Psychiatry*, 37, 896–901.
- 577 Myrtle, R.C., & Wilber, K.H. (1994). Designing service delivery systems: Lessons from the development of community-based systems of care for the elderly. *Public Administration Review*, 54(3), 245–252.

- 578 Reynolds, K., & Koster, V. (2005). *Innovative ways to finance mental health services in a primary care setting*. Washtenaw, MI: Washtenaw Community Health Association.
- 579 Davidson, L., Tondora, J., O'Connell, M.J., Kirk, T. Jr., Rockholz, P., & Evans, A.C. (2007). Creating a recovery-oriented system of behavioral health care: moving from concept to reality. *Psychiatric Rehabilitation Journal*, 31(1), 23–31.
- 580 Ibid.
- 581 O'Connell, M., Tondora, J., Croog, G., Evans, A., & Davidson, L. (2005).
- 582 Ibid.
- 583 Davidson, L., Tondora, J., O'Connell, M.J., et al. (2007).
- 584 White, W.L. (2007). A recovery revolution in Philadelphia. *Counselor*, 8(5), 34–38.
- 585 White, W.L. (2007). p. 36.
- 586 Ibid.
- 587 Ibid.
- 588 Ibid.
- 589 Barreira, P., Espey, B., Fishbein, R., Moran, D., & Flannery, R.B. (2000). Linking substance abuse and serious mental illness service delivery systems: Initiating a statewide collaborative. *Journal of Behavioral Health Services & Research*, 27(1), 107-113.
- 590 Barreira, P., Espey, B., Fishbein, R., et al. (2000).
- 591 Jacobson, N., & Curtis, L. (2000).
- 592 Solomon, P., & Stanhope, V. (2004).
- 593 Ibid. p. 318.
- 594 Ibid. p. 319.

